arXiv:1202.5728v1 [math.CA] 26 Feb 2012

A TRANSFERENCE RESULT OF THE L^p CONTINUITY OF THE JACOBI RIESZ TRANSFORM TO THE GAUSSIAN AND LAGUERRE RIESZ TRANSFORMS.

EDUARD NAVAS AND WILFREDO O. URBINA

Abstract. In this paper using the well known asymptotic relations between Jacobi polynomials and Hermite and Laguerre polynomials. we develop a transference method to obtain the L^p-continuity of the Gaussian-Riesz transform and the L^p-continuity of the Laguerre-Riesz transform from the L^p-continuity of the Jacobi-Riesz transform, in dimension one.

Dedicated to professor Calixto P. Calderón with deep admiration.

1. Preliminaries

In the theory of classical orthogonal polynomials, it is well know the asymptotic relations between Jacobi polynomials and Hermite and Laguerre polynomials. Base in those asymptotic relations we develop a transference method to obtain L^p-continuity for the Gaussian-Riesz transform and the L^p-continuity of the Laguerre-Riesz transform from the L^p-continuity of the Jacobi-Riesz transform, in dimension one.
For all the classical polynomials we are going to use the normalizations given in G. Segö's book [22].

- *Jacobi polynomials:* For $\alpha, \beta > -1$, the Jacobi polynomials $\{P_n^{(\alpha,\beta)}\}_{n\in\mathbb{N}}$ are defined as the orthogonal polynomials associated with the Jacobi measure $\mu_{\alpha,\beta}$ (or beta measure) in $(-1,1)$, defined as

$$\begin{aligned}\mu_{\alpha,\beta}(dx) &= \omega_{\alpha,\beta}(x)dx = \chi_{(-1,1)}(x)\frac{(1-x)^{\alpha}(1+x)^{\beta}}{2^{\alpha+\beta+1}B(\alpha+1,\beta+1)}dx \\ &= \eta_{\alpha,\beta}\chi_{(-1,1)}(x)(1-x)^{\alpha}(1+x)^{\beta}dx,\end{aligned} \tag{1.1}$$

where $\eta_{\alpha,\beta} = \frac{1}{2^{\alpha+\beta+1}B(\alpha+1,\beta+1)} = \frac{\Gamma(\alpha+\beta+2)}{2^{\alpha+\beta+1}\Gamma(\alpha+1)\Gamma(\beta+1)}$.

The function $\omega_{\alpha,\beta}$ is called the (normalized) *Jacobi weight.*

The Jacobi polynomials can be obtained from the canonical basis of the polynomials $\{1, x, x^2, \cdots, x^n, \cdots\}$ using the Gram-Schmidt orthogonalization process with respect to the inner product in $L^2(\mu_{\alpha,\beta})$. Thus we have the *orthogonality property* of Jacobi polynomials with respect to $\mu_{\alpha,\beta}$,

$$\int_{-\infty}^{\infty} P_n^{(\alpha,\beta)}(y)P_m^{(\alpha,\beta)}(y)\,\mu_{\alpha,\beta}(dy) = \eta_{\alpha,\beta}h_n^{(\alpha,\beta)}\delta_{n,m} = \hat{h}_n^{(\alpha,\beta)}\delta_{n,m}, \tag{1.2}$$

2000 Mathematics Subject Classification Primary 42C10; Secondary 26A24.
Key words and phrases: Transference, Riesz Transform, Orthogonal polynomials.

$n, m = 0, 1, 2, \cdots$, where

$$h_n^{(\alpha,\beta)} = \frac{2^{\alpha+\beta+1}}{(2n+\alpha+\beta+1)} \frac{\Gamma(n+\alpha+1)\,\Gamma(n+\beta+1)}{\Gamma(n+1)\,\Gamma(n+\alpha+\beta+1)}, \tag{1.3}$$

and

$$\begin{aligned} \hat{h_n}^{(\alpha,\beta)} &= \frac{1}{(2n+\alpha+\beta+1)} \frac{\Gamma(\alpha+\beta+2)\Gamma(n+\alpha+1)\,\Gamma(n+\beta+1)}{\Gamma(\alpha+1)\Gamma(\beta+1)\Gamma(n+1)\,\Gamma(n+\alpha+\beta+1)} \\ &= \|P_n^{(\alpha,\beta)}\|^2_{2,(\alpha,\beta)}. \end{aligned}$$

Moreover,

$$\begin{aligned} &\left\|P_{n-1}^{(\alpha+1,\beta+1)}\right\|^2_{2,(\alpha+1,\beta+1)} \\ = &\ \frac{1}{(2n+\alpha+\beta+1)} \frac{\Gamma(\alpha+\beta+4)\Gamma(n+\alpha+1)\,\Gamma(n+\beta+1)}{\Gamma(\alpha+2)\Gamma(\beta+2)\Gamma(n)\,\Gamma(n+\alpha+\beta+2)} \\ = &\ \frac{(\alpha+\beta+3)(\alpha+\beta+2)n}{(\alpha+1)(\beta+1)\,(n+\alpha+\beta+1)} \\ &\times \frac{\Gamma(\alpha+\beta+2)\Gamma(n+\alpha+1)\,\Gamma(n+\beta+1)}{(2n+\alpha+\beta+1)\Gamma(\alpha+1)\Gamma(\beta+1)\Gamma(n+1)\,\Gamma(n+\alpha+\beta+1)} \qquad (1.4) \\ = &\ \frac{(\alpha+\beta+3)(\alpha+\beta+2)n}{(\alpha+1)(\beta+1)\,(n+\alpha+\beta+1)} \left\|P_n^{(\alpha,\beta)}\right\|^2_{2,(\alpha,\beta)}. \end{aligned}$$

Now using the generalized Rodrigues' formula, Szegö [22],(4.10.1), for $m = 1$, we get

(1.5)
$$\frac{d}{dx}\left\{(1-x)^{\alpha+1}(1+x)^{\beta+1}P_{n-1}^{(\alpha+1,\beta+1)}(x)\right\} = -2n(1-x)^{\alpha}(1+x)^{\beta}P_n^{(\alpha,\beta)}(x).$$

On the other hand, the Jacobi polynomial $P_n^{(\alpha,\beta)}$ is a polynomial solution of the *Jacobi differential equation*, with parameters α, β, n,

$$\left(1-x^2\right)y'' + \left[\beta - \alpha - (\alpha+\beta+2)\,x\right]y' + n\,(n+\alpha+\beta+1)\,y = 0, \tag{1.6}$$

i. e. $P_n^{(\alpha,\beta)}$ is an eigenfunction of the (one-dimensional) second order diffusion operator

$$\mathcal{L}^{\alpha,\beta} = (1-x^2)\frac{d^2}{dx^2} + (\beta - \alpha - (\alpha+\beta+2)x)\frac{d}{dx}, \tag{1.7}$$

associated with the eigenvalue $-\lambda_n^{\alpha+\beta} = n(n+\alpha+\beta+1)$. $\mathcal{L}^{\alpha,\beta}$ is called the *Jacobi differential operator* .

The operator semigroup associated to the Jacobi polynomials is defined for positive or bounded measurable Borel functions of $(-1,1)$, as

$$T_t^{\alpha,\beta}f(x) = \int_{-1}^{1} p^{\alpha,\beta}(t,x,y)f(y)\mu_{\alpha,\beta}(dy), \tag{1.8}$$

where

$$p^{\alpha,\beta}(t,x,y) = \sum_k \frac{e^{-k(k+\alpha+\beta+1)t}}{\hat{h_k}^{(\alpha,\beta)}} P_k^{(\alpha,\beta)}(x)P_k^{(\alpha,\beta)}(y).$$

The explicit representation of $p^{\alpha,\beta}(t,x,y)$ is very complicated since the eigenvalues λ_n are not linearly distributed and was obtained by G. Gasper [5]. It is an analog of Bailey's F_4 representation for what is usually called the Jacobi-Poisson integral, see [2]. Considering the case From that form, taking $x=-y=1$, it can be proved that $p^{\alpha,\beta}(t,x,y)$ is a positive kernel for any $x\in(-1,1)$.

$\{T_t^{\alpha,\beta}\}$ is called the *Jacobi semigroup* and can be proved that is a Markov semigroup, for details see [24].

The Jacobi-Poisson semigroup $\{P_t^{\alpha,\beta}\}$ can be defined, using Bochner's subordination formula,

$$e^{-\lambda^{1/2}t}=\frac{1}{\sqrt{\pi}}\int_0^\infty \frac{e^{-u}}{\sqrt{u}}e^{-\frac{\lambda t^2}{4u}}du.$$

as the subordinated semigroup of the Jacobi semigroup,

$$P_t^{\alpha,\beta}f(x)=\frac{1}{\sqrt{\pi}}\int_0^\infty \frac{e^{-u}}{\sqrt{u}}T^{\alpha,\beta}_{t^2/4u}f(x)du.$$

For a function $f\in L^2\left([-1,1],\mu_{(\alpha,\beta)}\right)$ let us consider its Fourier-Jacobi expansion

$$f=\sum_{k=0}^\infty \frac{\langle f,P_k^{(\alpha,\beta)}\rangle}{\hat{h_k}^{(\alpha,\beta)}}P_k^{(\alpha,\beta)}, \tag{1.9}$$

where

$$\langle f,P_k^{(\alpha,\beta)}\rangle=\int_{-1}^1 f(y)P_k^{(\alpha,\beta)}(y)\mu_{\alpha,\beta}(dy).$$

Then, the action of T_t and P_t can be expressed as

$$T_t^{\alpha,\beta}f=\sum_{k=0}^\infty \frac{\langle f,P_k^{(\alpha,\beta)}\rangle}{\hat{h_k}^{(\alpha,\beta)}}e^{-\lambda_k t}P_k^{(\alpha,\beta)},$$

and

$$P_t^{\alpha,\beta}f=\sum_{k=0}^\infty \frac{\langle f,P_k^{(\alpha,\beta)}\rangle}{\hat{h_k}^{(\alpha,\beta)}}e^{-\sqrt{\lambda_k}t}P_k^{(\alpha,\beta)}.$$

Following the classical case, see [16], the *Jacobi-Riesz transform* can be define formally as

$$R^{\alpha,\beta}=\sqrt{1-x^2}\frac{d}{dx}(\mathcal{L}^{\alpha,\beta})^{-1/2}, \tag{1.10}$$

where $(\mathcal{L}^{\alpha,\beta})^{-\nu/2}$ is the Jacobi-Riesz potential of order $\nu/2$. $(\mathcal{L}^{\alpha,\beta})^{-\nu/2}$ can be represented as

$$(\mathcal{L}^{\alpha,\beta})^{-\nu/2}f=\frac{1}{\Gamma(\nu)}\int_0^\infty t^{\nu-1}P_t^{(\alpha,\beta)}f dt,$$

and then for $f\in L^2\left([-1,1],\mu_{(\alpha,\beta)}\right)$, $(\mathcal{L}^{\alpha,\beta})^{-\nu/2}f$ has Jacobi expansion

$$(\mathcal{L}^{\alpha,\beta})^{-\nu/2}f(x)=\sum_{k=0}^\infty \frac{\langle f,P_k^{(\alpha,\beta)}\rangle}{\hat{h_k}^{(\alpha,\beta)}}\lambda_k^{-\nu/2}P_k^{(\alpha,\beta)}(x). \tag{1.11}$$

Now, since

$$\frac{d}{dx}\left\{P_k^{(\alpha,\beta)}(x)\right\} = \frac{(k+\alpha+\beta+1)}{2}P_{k-1}^{(\alpha+1,\beta+1)}(x),$$

see Szegö [22],(4.21.7), and using (1.11) we get that the Jacobi-Riesz transform for $f \in L^2\left([-1,1], \mu_{(\alpha,\beta)}\right)$ has Jacobi expansion

$$R^{\alpha,\beta}f(x) = \sum_{k=1}^{\infty}\frac{\langle f, P_k^{(\alpha,\beta)}\rangle}{\hat{h}_k^{(\alpha,\beta)}}\lambda_k^{-1/2}\frac{(k+\alpha+\beta+1)}{2}\sqrt{1-x^2}P_{k-1}^{(\alpha+1,\beta+1)}(x), \tag{1.12}$$

where $\lambda_k = k(k+\alpha+\beta+1)$.

In particular, the *Gegenbauer polynomials* $\{C_n^\lambda\}$, $\lambda > -1/2$ are defined as

$$C_n^\lambda(x) = \frac{\Gamma(\lambda+1/2)\Gamma(n+2\lambda)}{\Gamma(2\lambda)\Gamma(n+\lambda+1/2)}P_n^{(\lambda-1/2,\lambda-1/2)}(x). \tag{1.13}$$

Gegenbauer polynomials are orthogonal with respect to the Gegenbauer measure, μ_λ in $(-1,1)$, defined as,

$$\begin{aligned}\mu_\lambda(dx) &= \omega_\lambda(x)dx = \chi_{(-1,1)}(x)\frac{\Gamma(2\lambda+1)}{2^{2\lambda}\left[\Gamma(\lambda+1/2)\right]^2}(1-x)^{\lambda-1/2}(1+x)^{\lambda-1/2}dx \\ &= \chi_{(-1,1)}(x)\frac{\lambda 2^{2\lambda}\left[\Gamma(\lambda)\right]^2}{2\pi\Gamma(2\lambda)}(1-x^2)^{\lambda-1/2}dx, \qquad (1.14)\end{aligned}$$

by Legendre's duplication formula, see [1] page 22.

Taking $\alpha = \beta = \lambda - 1/2$ in (1.4), then from (1.13) we have,

$$\begin{aligned}\|C_n^\lambda\|_{2,\lambda}^2 &= \frac{\left[\Gamma(\lambda+1/2)\right]^2\left[\Gamma(n+2\lambda)\right]^2}{\left[\Gamma(2\lambda)\right]^2\left[\Gamma(n+\lambda+1/2)\right]^2}\frac{\Gamma(2\lambda+1)\left[\Gamma(n+\lambda+1/2)\right]^2}{(2n+2\lambda)\left[\Gamma(\lambda+1/2)\right]^2\Gamma(n+1)\Gamma(n+2\lambda)} \\ &= \frac{\Gamma(2\lambda+1)\Gamma(n+2\lambda)}{\left[\Gamma(2\lambda)\right]^2 2(n+\lambda)n!} = \frac{\lambda\Gamma(n+2\lambda)}{\Gamma(2\lambda)(n+\lambda)n!}.\end{aligned} \tag{1.15}$$

Taking $\alpha = \beta = \lambda - 1/2$, in (1.5), we get

$$\begin{aligned}&\frac{d}{dx}\left\{(1-x^2)^{\lambda+1/2}C_{n-1}^{\lambda+1}(x)\right\} \\ &= -2n\frac{\Gamma(\lambda+3/2)\Gamma(n+2\lambda+1)}{\Gamma(2\lambda+2)\Gamma(n+\lambda+1/2)}\frac{\Gamma(2\lambda)\Gamma(n+\lambda+1/2)}{\Gamma(\lambda+1/2)\Gamma(n+2\lambda)}(1-x^2)^{\lambda-1/2}C_n^\lambda(x) \\ &= -n\frac{(2\lambda+1)(n+2\lambda)}{(2\lambda+1)(2\lambda)}(1-x^2)^{\lambda-1/2}C_n^\lambda(x) = -n\frac{(n+2\lambda)}{2\lambda}(1-x^2)^{\lambda-1/2}C_n^\lambda(x).\end{aligned} \tag{1.16}$$

Now, for $f \in L^2([-1,1], \mu_\lambda)$, from (1.12) we get that its *Gegenbauer-Riesz transform* will have the expansion

$$R^\lambda f(x) = \frac{1}{2}\sum_{k=1}^{\infty}\langle f, C_k^\lambda\rangle d_k^{(\lambda)}\left(\frac{k+2\lambda}{k}\right)^{1/2}\sqrt{1-x^2}C_{k-1}^{\lambda+1}(x), \tag{1.17}$$

where $d_k^{(\lambda)} = \frac{4\Gamma(2\lambda)(k+\lambda)\Gamma(k+1)}{\Gamma(k+2\lambda+1)}$.

The L^p-continuity of the Riesz-Jacobi transform $R^{\alpha,\beta}$, was proved by Zh. Li [8] and L. Caffarelli [4] in the case $d = 1$. In the case $d \geq 1$ $R_i^{\alpha,\beta}$, $i =, \cdots, d$, are defined analogously, using partial differentiation in (1.10), and their L^p-continuity was proved by A. Nowak and P. Sjogren, [15] Theorem 5.1.

Theorem 1.1. *Assume that* $1 < p < \infty$ *and* $\alpha, \beta \in [-1/2, \infty)^d$. *There exists a constant* c_p *such that*

$$\|R_i^{\alpha,\beta} f\|_{p,(\alpha,\beta)} \leq c_p \|f\|_{p,(\alpha,\beta)}. \tag{1.18}$$

for all $i = 1, \cdots, d$.

For the particular case of the Gegenbauer polynomials this result was obtained in the one dimensional case by B. Muckenhoupt and E. Stein in their seminal paper of 1965, [13].

- *Hermite polynomials*: The *Hermite polynomials* $\{H_n\}_n$, are defined as the orthogonal polynomials associated with the Gaussian measure in $\mathbb{R}$, $\gamma(dx) = \frac{e^{-x^2}}{\sqrt{\pi}} dx$, i.e.

$$\int_{-\infty}^{\infty} H_n(y) H_m(y)\, \gamma(dy) = 2^n n! \delta_{n,m}, \tag{1.19}$$

$n, m = 0, 1, 2, \cdots$, with the *normalization*

$$H_{2n+1}(0) = 0, \quad H_{2n}(0) = (-1)^n \frac{(2n)!}{n!}. \tag{1.20}$$

We have

$$H_n^{'}(x) \;=\; 2n H_{n-1}(x), \tag{1.21}$$

$$H_n^{''}(x) \;-\; 2x H_n^{'}(x) + 2n H_n(x) = 0, \tag{1.22}$$

thus H_n is an eigenfunction of the one dimensional *Ornstein-Uhlenbeck operator* or harmonic oscillator operator,

$$L = \frac{1}{2}\frac{d^2}{dx^2} - x\frac{d}{dx}, \tag{1.23}$$

associated with the eigenvalue $\lambda_n = -n$.

The *Gaussian-Riesz transform* can be defined formally, see [16], as

$$R^\gamma = \frac{d}{dx}(-L)^{-1/2}. \tag{1.24}$$

Therefore, for $f \in L^2(\mathbb{R}, \gamma)$ with Hermite expansion

$$f = \sum_{k=1}^{\infty} \frac{\langle f, H_k\rangle}{2^k k!} H_k,$$

its Gaussian-Riesz transform has Hermite expansion

$$R^\gamma f(x) = \sum_{k=1}^{\infty} \frac{\langle f, H_k\rangle}{2^k k!} \sqrt{2k} H_{k-1}(x). \tag{1.25}$$

The L^p continuity of the of the Gaussian-Riesz transform was proved by B. Muckenhoupt [11] in 1969, in the case $d = 1$. In the case $d \geq 1$ R_i^γ, $i =, \cdots, d$, are defined analogously, using partial differentiation in (1.24), and their L^p continuity has been proved by very different ways, using analytic and probabilistic tools, by P. A. Meyer [10], R. Gundy [6], S. Pérez and F. Soria [17], G. Pissier [18], C. Gutiérrez [7] and W. Urbina [23].

Theorem 1.2. *Assume that* $1 < p < \infty$*. There exists a constant* c_p *such that*

$$\|R_i^\gamma f\|_{p,\gamma} \leq c_p \|f\|_{p,\gamma}. \tag{1.26}$$

for all $i = 1, \cdots, d$.

- *Laguerre polynomials*: For $\alpha > -1$, the *Laguerre polynomials* $\{L_k^\alpha\}$ are defined as the orthogonal polynomials associated with the Gamma measure on $(0,\infty)$, $\mu_\alpha(dx) = \chi_{(0,\infty)}(x)\frac{x^\alpha e^{-x}}{\Gamma(\alpha+1)}dx$, i.e.

$$\int_0^\infty L_n^\alpha(y) L_m^\alpha(y)\, \mu_\alpha(dy) = \binom{n+\alpha}{n}\delta_{n,m} = \frac{\Gamma(n+\alpha+1)}{\Gamma(\alpha+1)n!}\delta_{n,m}, \tag{1.27}$$

$n, m = 0, 1, 2, \cdots$ We have

$$(L_k^\alpha(x))' = -L_{k-1}^{\alpha+1}(x). \tag{1.28}$$

$$x(L_k^\alpha(x))'' + (\alpha+1-x)(L_k^\alpha(x))' + kL_k^\alpha(x) = 0. \tag{1.29}$$

thus L_k^α is an eigenfunction of the (one-dimensional) *Laguerre differential operator*

$$\mathcal{L}^\alpha = x\frac{d^2}{dx^2} + (\alpha+1-x)\frac{d}{dx},$$

associated with the eigenvalue $\lambda_k = -k$

The *Laguerre-Riesz transform* can be defined formally, see [16], as

$$R^\alpha = \sqrt{x}\frac{d}{dx}(\mathcal{L}^\alpha)^{-1/2}. \tag{1.30}$$

Therefore for $f \in L^2((0,\infty), \mu_\alpha)$ with Laguerre expansion

$$f = \sum_{k=0}^\infty \frac{\Gamma(\alpha+1)k!}{\Gamma(k+\alpha+1)}\langle f, L_k^\alpha\rangle L_k^\alpha$$

its Laguerre-Riesz transform has Laguerre expansion

$$R^\alpha f(x) = -\sum_{k=1}^\infty \frac{\Gamma(\alpha+1)k!}{\Gamma(k+\alpha+1)}(\sqrt{k})^{-1}\sqrt{x}\langle f, L_k^\alpha\rangle L_{k-1}^{\alpha+1}(x). \tag{1.31}$$

The L^p continuity of the Laguerre-Riesz transform was proved by B. Muckenhoupt [12], for the case $d = 1$. In the case $d \geq 1$ R_i^α, $i =, \cdots, d$, are defined analogously, using partial differentiation in (1.30), and their L^p continuity was proved by A. Nowak [14] using Littlewood-Paley theory.

Theorem 1.3. *Assume that* $1 < p < \infty$ *and* $\alpha \in [-1/2, \infty)^d$*. There exists a constant* c_p *such that*

$$\|R_i^\alpha f\|_{p,\alpha} \leq c_p \|f\|_{p,\alpha}. \tag{1.32}$$

for all $i = 1, \cdots, d,$.

- Finally, the *asymptotic relations* between Jacobi polynomials and other classical orthogonal polynomials (see [22], (5.3.4) and (5.6.3)) are the following

i) With Hermite polynomials,

$$\lim_{\lambda\to\infty} \lambda^{-n/2} C_n^{\lambda}(x/\sqrt{\lambda}) = \frac{H_n(x)}{n!}, \tag{1.33}$$

ii) With Laguerre polynomials,

$$\lim_{\beta\to\infty} P_n^{(\alpha,\beta)}(1-2x/\beta) = L_n^{\alpha}(x). \tag{1.34}$$

Both relations holds uniformly in every closed interval of $\mathbb{R}$.

We want to thank Professor Luis A. Caffarelli to share with us the idea that such transferences between the Jacobi case and the Hermite and Laguerre cases should be possible.

2. MAIN RESULTS

We we want to obtain the L^p-continuity for the Gaussian-Riesz transform and for the L^p-continuity of the Laguerre-Riesz transform from the L^p-continuity of the Jacobi-Riesz transform, using a transference method based on the asymptotic relations between Jacobi polynomials and Hermite and Laguerre polynomials.
We will start considering the case $p = 2$; more precisely we want to prove

Theorem 2.1. *The $L^2(\mu_{\alpha,\beta})$ boundedness for the Jacobi-Riesz transform*

$$\|R^{\alpha,\beta} f\|_{2,(\alpha,\beta)} \leq C_2 \|f\|_{2,(\alpha,\beta)} \tag{2.1}$$

implies

i) *the $L^2(\gamma)$ boundedness for the Gaussian-Riesz transform*

$$\|R^{\gamma} f\|_{2,\gamma} \leq C_2 \|f\|_{2,\gamma}. \tag{2.2}$$

and

ii) *the $L^2(\mu_\alpha)$ boundedness for the Laguerre-Riesz transform*

$$\|R^{\alpha} f\|_{2,\alpha} \leq C_2 \|f\|_{2,\alpha}. \tag{2.3}$$

For the proof of this we will need the following technical result,

Proposition 2.1. *(norm relations)*

i) *Let $f \in L^2(\mathbb{R},\gamma)$ and define $f_\lambda(x) = f(\sqrt{\lambda}x) 1_{[-1,1]}(x)$, then $f_\lambda \in L^2([-1,1],\mu_\lambda)$ and*

$$\lim_{\lambda\to\infty} \|f_\lambda\|_{2,\lambda} = \|f\|_{2,\gamma} \tag{2.4}$$

ii) *Let $f \in L^2(\mathbb{R},\mu_\alpha)$ and define $f_\beta(x) = f\left(\frac{\beta}{2}(1-x)\right) 1_{[-1,1]}(x)$, then $f_\beta \in L^2([-1,1],\mu_{(\alpha,\beta)})$ and*

$$\lim_{\beta\to\infty} \|f_\beta\|_{2,(\alpha,\beta)} = \|f\|_{2,\alpha} \tag{2.5}$$

Proof

i) First let us prove that $f_\lambda \in L^2([-1,1],\mu_\lambda)$. For $x \in [-1,1]$,

$$(1-x^2)^{\lambda-1/2} \leq e^{-\lambda x^2} e^{\frac{x^2}{2}} \leq e^{-\lambda x^2} e^{\frac{1}{2}}. \tag{2.6}$$

Then,

$$\frac{\Gamma(2\lambda+1)}{2^{2\lambda}\left[\Gamma\left(\lambda+1/2\right)\right]^2}\int_{-1}^{1}(1-x^2)^{\lambda-1/2}|f_\lambda(x)|^2dx$$
$$\leq e^{\frac{1}{2}}\frac{\Gamma(2\lambda+1)}{2^{2\lambda}\left[\Gamma\left(\lambda+1/2\right)\right]^2}\int_{-1}^{1}|f(\sqrt{\lambda}x)|^2e^{-\lambda x^2}dx.$$

Now, making the change of variable $u=\sqrt{\lambda}x$, we have

$$\begin{aligned} e^{\frac{1}{2}}&\frac{\Gamma(2\lambda+1)}{2^{2\lambda}\left[\Gamma\left(\lambda+1/2\right)\right]^2}\int_{-1}^{1}|f(\sqrt{\lambda}x)|^2e^{-\lambda x^2}dx \\ &= e^{\frac{1}{2}}\frac{\Gamma(2\lambda+1)}{\sqrt{\lambda}2^{2\lambda}\left[\Gamma\left(\lambda+1/2\right)\right]^2}\int_{-\infty}^{\infty}|f(u)|^2 1_{[-\sqrt{\lambda},\sqrt{\lambda}]}e^{-u^2}du \\ &\leq e^{\frac{1}{2}}\frac{\Gamma(2\lambda+1)}{\sqrt{\lambda}2^{2\lambda}\left[\Gamma\left(\lambda+1/2\right)\right]^2}\int_{-\infty}^{\infty}|f(u)|^2e^{-u^2}du \\ &= e^{\frac{1}{2}}\frac{\Gamma(2\lambda+1)\sqrt{\pi}}{\sqrt{\lambda}2^{2\lambda}\left[\Gamma\left(\lambda+1/2\right)\right]^2}\|f\|_{2,\gamma}^2<\infty, \end{aligned}$$

as $f\in L^2(\mathbb{R},\gamma)$. Therefore $f_\lambda\in L^2([-1,1],\mu_\lambda)$.
On the other hand, using Legendre's duplication formula,

$$\begin{aligned} \|f_\lambda\|_{2,\lambda}^2 &= \frac{\Gamma(2\lambda+1)}{2^{2\lambda}\left[\Gamma\left(\lambda+1/2\right)\right]^2}\int_{-1}^{1}(1-x^2)^{\lambda-1/2}|f_\lambda(x)|^2dx \\ &= \frac{\lambda\left[\Gamma(\lambda)\right]^2 2^{2\lambda}}{2\pi\Gamma\left(2\lambda\right)}\int_{-1}^{1}(1-x^2)^{\lambda-1/2}|f_\lambda(x)|^2dx. \end{aligned}$$

Making the change of variable $x=\frac{y}{\sqrt{\lambda}}$, we get

$$\|f_\lambda\|_{2,\lambda}^2 = \frac{\lambda\left[\Gamma(\lambda)\right]^2 2^{2\lambda}}{2\pi\Gamma\left(2\lambda\right)}\int_{-\sqrt{\lambda}}^{\sqrt{\lambda}}|f_\lambda(\frac{y}{\sqrt{\lambda}})|^2(1-\frac{y^2}{\lambda})^{\lambda-1/2}\frac{dy}{\sqrt{\lambda}}.$$

Now, observe that

$$\lim_{\lambda\to\infty}f_\lambda(\frac{y}{\sqrt{\lambda}})=\lim_{\lambda\to\infty}f(y)1_{[-\sqrt{\lambda},\sqrt{\lambda}]}(y)=f(y),$$

Set, for all $y\in\mathbb{R}$

$$g(y,\lambda)=1_{[-\sqrt{\lambda},\sqrt{\lambda}]}(y)\left(f_\lambda(\frac{y}{\sqrt{\lambda}})\right)^2(1-\frac{y^2}{\lambda})^{\lambda-1/2},$$

then clearly

$$\lim_{\lambda\to\infty}g(y,\lambda)=(f(y))^2e^{-y^2},$$

thus by (2.6) we get

$$\begin{aligned} |g(y,\lambda)| &= 1_{[-\sqrt{\lambda},\sqrt{\lambda}]}(y)\left(f_\lambda(\frac{y}{\sqrt{\lambda}})\right)^2(1-\frac{y^2}{\lambda})^{\lambda-1/2} \\ &\leq e^{\frac{1}{2}}\left(f_\lambda(\frac{y}{\sqrt{\lambda}})\right)^2e^{-y^2}=e^{\frac{1}{2}}\left(f(y)\right)^2e^{-y^2}1_{[-\sqrt{\lambda},\sqrt{\lambda}]}(y)<e^{\frac{1}{2}}\left(f(y)\right)^2e^{-y^2}. \end{aligned}$$

Then, for $f \in L^2(\mathbb{R},\gamma)$, by the dominated convergence theorem, we have

$$\begin{aligned}\lim_{\lambda\to\infty}\int_{-\infty}^{\infty}|f_\lambda(\frac{y}{\sqrt{\lambda}})|^2 1_{[-\sqrt{\lambda},\sqrt{\lambda}]}(y)(1-\frac{y^2}{\lambda})^{\lambda-1/2}dy &= \int_{-\infty}^{\infty}\lim_{\lambda\to\infty} g(y,\lambda)dy\\ &= \int_{-\infty}^{\infty}(f(y))^2 e^{-y^2}dy.\end{aligned}$$

On the other hand, using the identity

$$\lim_{z\to\infty} z^{b-a}\frac{\Gamma(z+a)}{\Gamma(z+b)} = 1$$

and Legendre's duplication formula, we get,

$$\begin{aligned}\lim_{\lambda\to\infty}\frac{\lambda\,[\Gamma(\lambda)]^2\,2^{2\lambda}}{\lambda^{1/2}2\pi\Gamma\,(2\lambda)} &= \lim_{\lambda\to\infty}\frac{\lambda\,[\Gamma(\lambda)]^2\,2^{2\lambda}}{2\pi(2\pi)^{-1/2}2^{2\lambda-1/2}\Gamma\,(\lambda)\,\Gamma\left(\lambda+\frac{1}{2}\right)\lambda^{1/2}}\\ &= \lim_{\lambda\to\infty}\frac{\lambda\Gamma(\lambda)}{\lambda^{1/2}\sqrt{\pi}\Gamma\left(\lambda+\frac{1}{2}\right)} = \lim_{\lambda\to\infty}\frac{\lambda^{-1/2}\Gamma(\lambda+1)}{\sqrt{\pi}\Gamma\left(\lambda+\frac{1}{2}\right)} = \frac{1}{\sqrt{\pi}}.\end{aligned}$$

Then,

$$\begin{aligned}\lim_{\lambda\to\infty}\|f_\lambda\|^2_{2,\lambda} &= \lim_{\lambda\to\infty}\frac{\lambda\,[\Gamma(\lambda)]^2\,2^{2\lambda}}{2\pi\Gamma\,(2\lambda)}\int_{-\sqrt{\lambda}}^{\sqrt{\lambda}}|f_\lambda(\frac{y}{\sqrt{\lambda}})|^2(1-\frac{y^2}{\lambda})^{\lambda-1/2}\frac{dy}{\sqrt{\lambda}}\\ &= \frac{1}{\sqrt{\pi}}\int_{-\infty}^{\infty}|f(y)|^2 e^{-y^2}dy = \|f\|^2_{2,\gamma}.\end{aligned}$$

ii) For the Laguerre case we have,

$$\begin{aligned}&\frac{1}{2^{\alpha+\beta+1}B(\alpha+1,\beta+1)}\int_{-1}^{1}(1-x)^\alpha(1+x)^\beta|f_\beta(x)|^2dx\\ &\qquad= \frac{\Gamma(\alpha+\beta+2)}{2^{\alpha+\beta+1}\Gamma(\alpha+1)\Gamma(\beta+1)}\int_{-1}^{1}(1-x)^\alpha(1+x)^\beta|f_\beta(x)|^2dx,\end{aligned}$$

and using the change of variable $x = 1-\frac{2y}{\beta}$,

$$\begin{aligned}&\frac{\Gamma(\alpha+\beta+2)}{2^{\alpha+\beta+1}\Gamma(\alpha+1)\Gamma(\beta+1)}\frac{2}{\beta}\int_0^\beta\left(\frac{2y}{\beta}\right)^\alpha\left(2-\frac{2y}{\beta}\right)^\beta|f_\beta(1-\frac{2y}{\beta})|^2dy\\ &\qquad= \frac{\Gamma(\alpha+\beta+2)}{\Gamma(\alpha+1)\Gamma(\beta+1)\beta^{\alpha+1}}\int_0^\beta y^\alpha\left(1-\frac{y}{\beta}\right)^\beta|f(y)|^2dy.\end{aligned}$$

Then

$$\begin{aligned}&\frac{\Gamma(\alpha+\beta+2)}{\Gamma(\alpha+1)\Gamma(\beta+1)\beta^{\alpha+1}}\int_0^\beta y^\alpha\left(1-\frac{y}{\beta}\right)^\beta|f(y)|^2dy\\ &\qquad\leq \frac{\Gamma(\alpha+\beta+2)}{\Gamma(\alpha+1)\Gamma(\beta+1)\beta^{\alpha+1}}\int_0^\infty y^\alpha e^{-y}|f(y)|^2dy\\ &\qquad= \frac{\Gamma(\alpha+\beta+2)}{\Gamma(\beta+1)\beta^{\alpha+1}}\|f\|^2_{2,\alpha} < \infty,\end{aligned}$$

as $f \in L^2(\mathbb{R},\mu_\alpha)$, and therefore $f_\beta \in L^2([-1,1],\mu_{(\alpha,\beta)})$.

On the other hand, making the change of variable $x = 1 - \frac{2y}{\beta}$

$$\begin{aligned}\|f_\beta\|^2_{2,(\alpha,\beta)} &= \frac{\Gamma(\alpha+\beta+2)}{2^{\alpha+\beta+1}\Gamma(\alpha+1)\Gamma(\beta+1)}\frac{2}{\beta}\int_0^\beta \left(\frac{2y}{\beta}\right)^\alpha \left(2-\frac{2y}{\beta}\right)^\beta |f_\beta(1-\frac{2y}{\beta})|^2 dy\\ &= \frac{\Gamma(\beta+(\alpha+2))}{\beta^{\alpha+2}\Gamma(\alpha+1)\Gamma(\beta)}\int_0^\infty y^\alpha \left(1-\frac{y}{\beta}\right)^\beta |f_\beta(1-\frac{2y}{\beta})|^2 1_{[0,\beta]}(y)dy.\end{aligned}$$

Now, for β big enough,

$$\frac{\Gamma(\beta+(\alpha+2))}{\Gamma(\beta)} \simeq \beta^{\alpha+2}, \tag{2.7}$$

and therefore

$$\|f_\beta\|^2_{2,(\alpha,\beta)} \simeq \frac{1}{\Gamma(\alpha+1)}\int_0^\infty y^\alpha \left(1-\frac{y}{\beta}\right)^\beta |f_\beta(1-\frac{2y}{\beta})|^2 1_{[0,\beta]}(y)dy,$$

for β big enough. Observe that,

$$\lim_{\beta\to\infty} f_\beta(1-\frac{2y}{\beta}) = \lim_{\beta\to\infty} f(y)1_{[0,\beta]}(y) = f(y),$$

so let

$$s(y,\beta) = y^\alpha \left(1-\frac{y}{\beta}\right)^\beta |f_\beta(1-\frac{2y}{\beta})|^2 1_{[0,\beta]}(y),$$

then

$$\lim_{\beta\to\infty} s(y,\beta) = y^\alpha e^{-y}|f(y)|^2,$$

but as

$$|s(y,\beta)| \leq y^\alpha e^{-y}|f_\beta(1-\frac{2y}{\beta})|^2 1_{[0,\beta]}(y) \leq y^\alpha e^{-y}|f(y)|^2,$$

then, by the dominated convergent theorem, we get

$$\begin{aligned}\lim_{\beta\to\infty}\int_0^\infty y^\alpha \left(1-\frac{y}{\beta}\right)^\beta |f_\beta(1-\frac{2y}{\beta})|^2 1_{[0,\beta]}(y)dy &= \lim_{\beta\to\infty}\int_0^\infty s(y,\beta)dy\\ &= \int_0^\infty y^\alpha e^{-y}|f(y)|^2 dy.\end{aligned}$$

Then

$$\lim_{\beta\to\infty}\|f_\beta\|^2_{2,(\alpha,\beta)} = \frac{1}{\Gamma(\alpha+1)}\int_0^\infty y^\alpha e^{-y}|f(y)|^2 dy = \|f\|^2_{2,\alpha}. \blacksquare$$

The next result is analogous to the previous one but refer to the inner product and it will be crucial in the proof of Theorem 2.1.

Proposition 2.2. *(inner product relations) With the same notation as in Lemma 2.1,*

i) *Let $f \in L^2(\mathbb{R},\gamma)$, then*

$$\lim_{\lambda\to\infty}\langle f_\lambda, \lambda^{-k/2}C_k^\lambda\rangle = \langle f, \frac{H_k}{k!}\rangle \tag{2.8}$$

ii) *Let $f \in L^2(\mathbb{R},\mu_\alpha)$, then*

$$\lim_{\beta\to\infty}\langle f_\beta, P_k^{(\alpha,\beta)}\rangle = \langle f, L_k^\alpha\rangle \tag{2.9}$$

Proof

i) Let us prove first that

$$\lim_{\lambda\to\infty}\left\|f_\lambda+\lambda^{-k/2}C_k^\lambda\right\|_{2,\lambda}^2=\left\|f+\frac{H_k}{k!}\right\|_{2,\gamma}^2. \tag{2.10}$$

In fact, given that,

$$\frac{\lambda^{1/2}\left[\Gamma(\lambda)\right]^2 2^{2\lambda}}{2\pi\Gamma\left(2\lambda\right)}(1-\frac{y^2}{\lambda})^{\lambda-1/2}\left(f_\lambda(\frac{y}{\sqrt{\lambda}})+\lambda^{-k/2}C_k^\lambda(\frac{y}{\sqrt{\lambda}})\right)^2 1_{[-\sqrt{\lambda},\sqrt{\lambda}]}\geq 0,$$

by Fatou's lemma we get

$$\begin{aligned}&\frac{1}{\sqrt{\pi}}\int_{-\infty}^{\infty}\left(f(y)+\frac{H_k}{k!}(y)\right)^2 e^{-y^2}dy\\ &\qquad\leq\lim_{\lambda\to\infty}\frac{\lambda^{1/2}\left[\Gamma(\lambda)\right]^2 2^{2\lambda}}{2\pi\Gamma\left(2\lambda\right)}\int_{-\sqrt{\lambda}}^{\sqrt{\lambda}}\left(f_\lambda(\frac{y}{\sqrt{\lambda}})+\lambda^{-k/2}C_k^\lambda(\frac{y}{\sqrt{\lambda}})\right)^2(1-\frac{y^2}{\lambda})^{\lambda-1/2}dy,\end{aligned}$$

thus

$$\left\|f+\frac{H_k}{k!}\right\|_{2,\gamma}^2\leq\lim_{\lambda\to\infty}\left\|f_\lambda+\lambda^{-k/2}C_k^\lambda\right\|_{2,\lambda}^2.$$

On the other hand, let

$$G(y,\lambda)=\left(\frac{\lambda^{1/2}\left[\Gamma(\lambda)\right]^2 2^{2\lambda}}{2\pi\Gamma\left(2\lambda\right)}\right)^{1/2}\left(f_\lambda(\frac{y}{\sqrt{\lambda}})+\lambda^{-k/2}C_k^\lambda(\frac{y}{\sqrt{\lambda}})\right)(1-\frac{y^2}{\lambda})^{-1/4}1_{[-\sqrt{\lambda},\sqrt{\lambda}]}(y).$$

Clearly $G\in L^2(\mathbb{R},\gamma)$, and moreover

$$\lim_{\lambda\to\infty}G(y,\lambda)=\frac{1}{\pi^{1/4}}\left(f(y)+\frac{H_k}{k!}(y)\right)^2.$$

Then,

$$\begin{aligned}&\left\|f_\lambda+\lambda^{-k/2}C_k^\lambda\right\|_{2,\lambda}^2\\ &=\int_{-\infty}^{\infty}\frac{\lambda^{1/2}\left[\Gamma(\lambda)\right]^2 2^{2\lambda}}{2\pi\Gamma\left(2\lambda\right)}\left(f_\lambda(\frac{y}{\sqrt{\lambda}})+\lambda^{-k/2}C_k^\lambda(\frac{y}{\sqrt{\lambda}})\right)^2(1-\frac{y^2}{\lambda})^{\lambda-1/2}1_{[-\sqrt{\lambda},\sqrt{\lambda}]}dy\\ &=\int_{-\infty}^{\infty}\left(G(y,\lambda)\right)^2(1-\frac{y^2}{\lambda})^{\lambda}dy\leq\int_{-\infty}^{\infty}\left(G(y,\lambda)\right)^2 e^{-y^2}dy=\sqrt{\pi}\left\|G(\cdot,\lambda)\right\|_{2,\gamma}^2.\end{aligned}$$

Then by the continuity of the L^2 norm

$$\begin{aligned}\lim_{\lambda\to\infty}\left\|f_\lambda+\lambda^{-k/2}C_k^\lambda\right\|_{2,\lambda}^2 &\leq \lim_{\lambda\to\infty}\sqrt{\pi}\left\|G(\cdot,\lambda)\right\|_{2,\gamma}^2=\sqrt{\pi}\left\|\lim_{\lambda\to\infty}G(\cdot,\lambda)\right\|_{2,\gamma}^2\\ &= \sqrt{\pi}\left\|\frac{1}{\pi^{1/4}}\left(f+\frac{H_k}{k!}\right)\right\|_{2,\gamma}^2=\left\|f+\frac{H_k}{k!}\right\|_{2,\gamma}^2.\end{aligned}$$

Therefore

$$\lim_{\lambda\to\infty}\left\|f_\lambda+\lambda^{-k/2}C_k^\lambda\right\|_{2,\lambda}^2=\left\|f+\frac{H_k}{k!}\right\|_{2,\gamma}^2,$$

and analogously,

$$\lim_{\lambda\to\infty}\left\|f_\lambda-\lambda^{-k/2}C_k^\lambda\right\|_{2,\lambda}^2=\left\|f-\frac{H_k}{k!}\right\|_{2,\gamma}^2.$$

Then using the polarization formula,

$$\begin{aligned}\lim_{\lambda\to\infty}\langle f_\lambda,\lambda^{-k/2}C_k^\lambda\rangle &= \lim_{\lambda\to\infty}\frac{1}{4}\left[\left\|f_\lambda+\lambda^{-k/2}C_k^\lambda\right\|_{2,\lambda}^2-\left\|f_\lambda-\lambda^{-k/2}C_k^\lambda\right\|_{2,\lambda}^2\right]\\ &= \frac{1}{4}\left[\left\|f+\frac{H_k}{k!}\right\|_{2,\gamma}^2-\left\|f-\frac{H_k}{k!}\right\|_{2,\gamma}^2\right]=\langle f,\frac{H_k}{k!}\rangle.\end{aligned}$$

ii) Analogously as in part i), let us prove first that

$$\lim_{\beta\to\infty}\left\|f_\beta+P_k^{(\alpha,\beta)}\right\|_{2,(\alpha,\beta)}^2=\|f+L_k^\alpha\|_{2,\alpha}^2. \tag{2.11}$$

By Fatou's lemma, we have

$$\begin{aligned}&\frac{1}{\Gamma(\alpha+1)}\int_0^\infty (f(y)+L_k^\alpha(y))^2\,y^\alpha e^{-y}dy,\\ &\le \lim_{\beta\to\infty}\frac{\Gamma(\beta+(\alpha+2))}{\beta^{\alpha+2}\Gamma(\alpha+1)\Gamma(\beta)}\int_0^\infty\left|f_\beta(1-\frac{2y}{\beta})+P_k^{(\alpha,\beta)}(1-\frac{2y}{\beta})\right|^2 1_{[0,\beta]}(y)y^\alpha\left(1-\frac{y}{\beta}\right)^\beta dy,\end{aligned}$$

Then,

$$\|f+L_k^\alpha\|_{2,\alpha}^2\le\lim_{\beta\to\infty}\left\|f_\beta+P_k^{(\alpha,\beta)}\right\|_{2,(\alpha,\beta)}^2.$$

On the other hand, set

$$S(y,\beta)=\left(\frac{\Gamma(\beta+(\alpha+2))}{\beta^{\alpha+2}\Gamma(\alpha+1)\Gamma(\beta)}\right)^{1/2}\left(f_\beta(1-\frac{2y}{\beta})+P_k^{(\alpha,\beta)}(1-\frac{2y}{\beta})\right)1_{[0,\beta]}(y).$$

It is clear that $S\in L^2(\mathbb{R},\mu_\alpha)$, and moreover

$$\lim_{\beta\to\infty}S(y,\beta)=\left(\frac{1}{\Gamma(\alpha+1)}\right)^{1/2}(f(y)+L_k^\alpha(y)).$$

Now,

$$\begin{aligned}&\left\|f_\beta+P_k^{(\alpha,\beta)}\right\|_{2,(\alpha,\beta)}^2\\ &=\int_0^\infty\frac{\Gamma(\beta+(\alpha+2))}{\beta^{\alpha+2}\Gamma(\alpha+1)\Gamma(\beta)}\left|f_\beta(1-\frac{2y}{\beta})+P_k^{(\alpha,\beta)}(1-\frac{2y}{\beta})\right|^2 1_{[0,\beta]}y^\alpha\left(1-\frac{y}{\beta}\right)^\beta dy\\ &=\int_0^\infty (S(y,\beta))^2y^\alpha\left(1-\frac{y}{\beta}\right)^\beta dy\le\int_{-\infty}^\infty (S(y,\beta))^2y^\alpha e^{-y}dy=\Gamma(\alpha+1)\|S(\cdot,\beta)\|_{2,\alpha}^2,\end{aligned}$$

and by the continuity of the L^2-norm

$$\lim_{\beta\to\infty}\Gamma(\alpha+1)\|S(\cdot,\beta)\|_{2,\alpha}^2=\Gamma(\alpha+1)\left\|\lim_{\beta\to\infty}S(\cdot,\beta)\right\|_{2,\alpha}^2.$$

Thus,

$$\begin{aligned}\lim_{\beta\to\infty}\left\|f_\beta+P_k^{(\alpha,\beta)}\right\|_{2,(\alpha,\beta)}^2 &\leq \lim_{\beta\to\infty}\Gamma(\alpha+1)\left\|S(\cdot,\beta)\right\|_{2,\alpha}^2=\Gamma(\alpha+1)\left\|\lim_{\beta\to\infty}S(\cdot,\beta)\right\|_{2,\alpha}^2\\ &= \Gamma(\alpha+1)\left\|S(\cdot,\beta)\right\|_{2,\alpha}^2=\|f+L_k^\alpha\|_{2,\alpha}^2.\end{aligned}$$

Therefore,

$$\lim_{\beta\to\infty}\left\|f_\beta+P_k^{(\alpha,\beta)}\right\|_{2,(\alpha,\beta)}^2=\|f+L_k^\alpha\|_{2,\alpha}^2,$$

and analogously,

$$\lim_{\beta\to\infty}\left\|f_\beta-P_k^{(\alpha,\beta)}\right\|_{2,(\alpha,\beta)}^2=\|f-L_k^\alpha\|_{2,\alpha}^2.$$

Then by the polarization formula,

$$\begin{aligned}\lim_{\beta\to\infty}\langle f_\beta,P_k^{(\alpha,\beta)}\rangle &= \lim_{\beta\to\infty}\frac{1}{4}\left[\left\|f_\beta+P_k^{(\alpha,\beta)}\right\|_{2,(\alpha,\beta)}^2-\left\|f_\beta-P_k^{(\alpha,\beta)}\right\|_{2,(\alpha,\beta)}^2\right]\\ &= \frac{1}{4}\left[\|f+L_k^\alpha\|_{2,\alpha}^2-\|f-L_k^\alpha\|_{2,\alpha}^2\right]=\langle f,L_k^\alpha\rangle. \quad \blacksquare\end{aligned}$$

We are ready to prove Theorem 2.1,

Proof

First of all let us observe that by Parseval's identity, for $f\in L^2\left([-1,1],\mu_\lambda\right)$

$$\left\|R^{(\alpha,\beta)}f\right\|_{2,(\alpha,\beta)}^2 = \sum_{k=1}^{\infty}\left|\frac{\langle f,P_k^{\alpha,\beta}\rangle}{\hat{h_n}^{(\alpha,\beta)}}\right|^2\frac{(k+\alpha+\beta+1)^2}{4\lambda_k}\left\|\sqrt{1-x^2}P_{k-1}^{(\alpha+1,\beta+1)}\right\|_{2,(\alpha,\beta)}^2.$$

Since,

$$\begin{aligned}&\left\|\sqrt{1-x^2}P_{k-1}^{(\alpha+1,\beta+1)}\right\|_{2,(\alpha,\beta)}^2\\ &=\frac{4(\alpha+1)(\beta+1)}{2^{\alpha+\beta+3}(\alpha+\beta+3)(\alpha+\beta+2)B(\alpha+2,\beta+2)}\\ &\qquad\times\int_{-1}^{1}(1-x)^{\alpha+1}(1+x)^{\beta+1}\left[P_{k-1}^{(\alpha+1,\beta+1)}(x)\right]^2dx\\ &=\frac{4(\alpha+1)(\beta+1)}{(\alpha+\beta+3)(\alpha+\beta+2)}\left\|P_{k-1}^{(\alpha+1,\beta+1)}\right\|_{2,(\alpha+1,\beta+1)}^2=\frac{4k}{(k+\alpha+\beta+1)}\left\|P_k^{(\alpha,\beta)}\right\|_{2,(\alpha,\beta)}^2,\end{aligned}$$

we get,

$$\begin{aligned}\left\|R^{\alpha,\beta}f\right\|_{2,(\alpha,\beta)}^2 &= \sum_{k=1}^{\infty}\left|\frac{\langle f,P_k^{(\alpha,\beta)}\rangle}{\hat{h_n}^{(\alpha,\beta)}}\right|^2\frac{(k+\alpha+\beta+1)^2}{4k(k+\alpha+\beta+1)}\frac{4k}{(k+\alpha+\beta+1)}\left\|P_k^{(\alpha,\beta)}\right\|_{2,(\alpha,\beta)}^2\\ &= \sum_{k=1}^{\infty}|\langle f,P_k^{(\alpha,\beta)}\rangle|^2\\ &\qquad\times\frac{(2k+\alpha+\beta+1)\Gamma(\alpha+1)\Gamma(\beta+1)\Gamma(k+1)\Gamma(k+\alpha+\beta+1)}{\Gamma(\alpha+\beta+2)\Gamma(k+\alpha+1)\Gamma(k+\beta+1)}\end{aligned}$$

i) Now, for the Gegenbauer case, $\alpha=\beta=\lambda-1/2$, we have

$$\begin{aligned}\left\|R^\lambda f\right\|_{2,\lambda}^2 &= \sum_{k=1}^{\infty}|\langle f,C_k^\lambda\rangle|^2\frac{(2k+2\lambda)\,[\Gamma(\lambda+1/2)]^2\,\Gamma(k+1)\Gamma(k+2\lambda)}{\Gamma(2\lambda+1)\,[\Gamma(k+\lambda+1/2)]^2}\\ &\qquad\qquad\times\frac{[\Gamma(2\lambda)]^2\,[\Gamma(k+\lambda+1/2)]^2}{[\Gamma(\lambda+1/2)]^2\,[\Gamma(k+2\lambda)]^2}\\ &= \sum_{k=1}^{\infty}|\langle f,C_k^\lambda\rangle|^2\frac{2(k+\lambda)\,[\Gamma(2\lambda)]^2\,\Gamma(k+1)}{\Gamma(2\lambda+1)\Gamma(k+\lambda+1/2)}\\ &= \sum_{k=1}^{\infty}|\langle f,C_k^\lambda\rangle|^2\frac{(k+\lambda)\Gamma(2\lambda)\Gamma(k+1)}{\lambda\Gamma(k+2\lambda)} = \sum_{k=1}^{\infty}|\langle f,C_k^\lambda\rangle|^2\frac{(k+\lambda)k!}{\lambda(2\lambda)_k}\\ &\geq \sum_{k=1}^{\infty}|\langle f,\lambda^{-k/2}C_k^\lambda\rangle|^2\frac{(\frac{k}{\lambda}+1)k!}{2\sqrt{\pi}(2+1/\lambda)(2+2/\lambda)\dots(2+(k-1)/\lambda)}.\end{aligned}$$

Then, we conclude that

$$\sum_{k=1}^{\infty}|\langle f,\lambda^{-k/2}C_k^\lambda\rangle|^2\frac{(\frac{k}{\lambda}+1)k!}{2\sqrt{\pi}(2+\frac{1}{\lambda})(2+\frac{2}{\lambda})\dots(2+\frac{(k-1)}{\lambda})}\leq\left\|R^\lambda f\right\|_{2,\lambda}^2.$$

On the other hand, again using Parseval's identity, we have that the L^2-norm of the Gaussian Riesz transform for $f\in L^2(\mathbb{R},\gamma)$ is given by

$$\|R^\gamma f\|_{2,\gamma}^2 = \sum_{k=1}^{\infty}\frac{|\langle f,H_k\rangle|^2}{k!2^k\sqrt{\pi}}.$$

Now, since $f_\lambda(x)=f(\sqrt{\lambda}x)1_{[-1,1]}(x)$ taking $\lambda\to\infty$ in (2.12) and using the asymptotic relation (1.33), we get

$$\begin{aligned}\|R^\gamma f\|_{2,\gamma}^2 &= \frac{1}{\sqrt{\pi}}\lim_{\lambda\to\infty}\sum_{k=1}^{\infty}\frac{(\frac{k}{\lambda}+1)k!}{2\sqrt{\pi}(2+\frac{1}{\lambda})(2+\frac{2}{\lambda})\dots(2+\frac{(k-1)}{\lambda})}|\langle f_\lambda,\lambda^{-k/2}C_k^\lambda\rangle|^2\\ &\leq \frac{1}{\sqrt{\pi}}\lim_{\lambda\to\infty}\left\|R^\lambda f_\lambda\right\|_{2,\lambda}^2.\end{aligned}$$

Therefore, using Theorem 1.1 and Proposition 2.1 i), we get

$$\|R^\gamma f\|_{2,\gamma}^2 \leq \lim_{\lambda\to\infty}\left\|R^\lambda f_\lambda\right\|_{2,\lambda}^2\leq C_2\lim_{\lambda\to\infty}\|f_\lambda\|_{2,\lambda}^2=C_2\,\|f\|_{2,\gamma}^2.$$

ii) Observe that

$$\|R^\alpha f\|_{2,\alpha}^2=\sum_{k=1}^{\infty}\frac{[\Gamma(\alpha+1)k!]^2|\langle f,L_k^\alpha\rangle|^2}{k[\Gamma(k+\alpha+1)]^2}\|\sqrt{x}L_{k-1}^{\alpha+1}\|_{2,\alpha}^2.$$

But since

$$\begin{aligned}\|\sqrt{x}L_{k-1}^{\alpha+1}\|_{2,\alpha}^2 &= \int_0^{\infty}[L_{k-1}^{\alpha+1}(x)]^2x^\alpha e^{-x}\frac{dx}{\Gamma(\alpha+1)}=\frac{\Gamma(\alpha+2)}{\Gamma(\alpha+1)}\frac{\Gamma(k+\alpha+1)}{(k-1)!\Gamma(\alpha+2)}\\ &= \frac{\Gamma(k+\alpha+1)}{(k-1)!\Gamma(\alpha+1)},\end{aligned}$$

then

$$\begin{aligned}\|R^{\alpha}f\|_{2,\alpha}^2 &= \sum_{k=1}^{\infty}\frac{[\Gamma(\alpha+1)k!]^2|\langle f,L_k^{\alpha}\rangle|^2}{k[\Gamma(k+\alpha+1)]^2}\frac{\Gamma(k+\alpha+1)}{(k-1)!\Gamma(\alpha+1)}\\ &= \sum_{k=1}^{\infty}\frac{\Gamma(\alpha+1)k!|\langle f,L_k^{\alpha}\rangle|^2}{\Gamma(k+\alpha+1)}.\end{aligned}$$

Therefore

$$\|R^{\alpha}f\|_{2,\alpha}^2 = \lim_{\beta\to\infty}\sum_{k=1}^{\infty}\frac{\Gamma(\alpha+1)k!|\langle f_\beta,P_k^{(\alpha,\beta)}\rangle|^2}{\Gamma(k+\alpha+1)},$$

where, as before $f_\beta(x) = f\left(\frac{\beta}{2}(1-x)\right)1_{[-1,1]}(x)$. Now, for β big enough

$$\frac{\Gamma(\beta+(k+\alpha+1))}{\Gamma(\beta+(k+1))\beta^{\alpha}} \simeq 1 \qquad \text{and} \qquad \frac{\Gamma(\beta+1)}{\Gamma(\beta+(\alpha+2))\beta^{-\alpha-1}} \simeq 1,$$

then

$$\begin{aligned}&\sum_{k=1}^{\infty}|\langle f_\beta,P_k^{(\alpha,\beta)}\rangle|^2\frac{\Gamma(\alpha+1)k!}{\Gamma(k+\alpha+1)}\\ &= \sum_{k=1}^{\infty}|\langle f_\beta,P_k^{(\alpha,\beta)}\rangle|^2\frac{\Gamma(\alpha+1)k!}{\Gamma(k+\alpha+1)}\frac{\Gamma(\beta+(k+\alpha+1))}{\Gamma(\beta+(k+1))\beta^{\alpha}}\frac{\Gamma(\beta+1)}{\Gamma(\beta+\alpha+2)\beta^{-\alpha-1}}\\ &\leq \sum_{k=1}^{\infty}|\langle f_\beta,P_k^{(\alpha,\beta)}\rangle|^2\frac{(2k+\alpha+\beta+1)k!\Gamma(\alpha+1)\Gamma(\beta+1)\Gamma(k+\beta+\alpha+1)}{\Gamma(\beta+\alpha+2)\Gamma(k+\alpha+1)\Gamma(k+\beta+1)},\end{aligned}$$

for β big enough. Thus

$$\sum_{k=1}^{\infty}|\langle f_\beta,P_k^{(\alpha,\beta)}\rangle|^2\frac{\Gamma(\alpha+1)k!}{\Gamma(k+\alpha+1)} \leq \left\|R^{(\alpha,\beta)}f_\beta\right\|_{2,(\alpha,\beta)}^2 \tag{2.12}$$

for β big enough. Now, taking $\beta\to\infty$ and using the asymptotic relation (1.34) in (2.12) we get,

$$\begin{aligned}\|R^{\alpha}f\|_{2,\alpha}^2 &= \lim_{\beta\to\infty}\sum_{k=1}^{\infty}\frac{\Gamma(\alpha+1)k!|\langle f_\beta,P_k^{(\alpha,\beta)}\rangle|^2}{\Gamma(k+\alpha+1)}\\ &\leq \lim_{\beta\to\infty}\left\|R^{(\alpha,\beta)}f_\beta\right\|_{2,(\alpha,\beta)}^2.\end{aligned}$$

Therefore, using Theorem 1.1 and Proposition 2.1 ii), we get

$$\|R^{\alpha}f\|_{2,\alpha}^2 \leq \lim_{\beta\to\infty}\left\|R^{(\alpha,\beta)}f_\beta\right\|_{2,(\alpha,\beta)}^2 \leq C_2\lim_{\lambda\to\infty}\|f_\beta\|_{2,(\alpha,\beta)}^2 = C_2\|f\|_{2,\alpha}^2. \blacksquare$$

Now we are going to consider the general case $p \neq 2$. For the proof we will follow the argument given by Betancour et al in [3].

Theorem 2.2. *Let $\alpha,\beta > -1$ and $1 < p < \infty$, then the $L^p(\mu_{\alpha,\beta})$ boundedness for the Jacobi-Riesz transform*

$$\|R^{\alpha,\beta}f\|_{p,(\alpha,\beta)} \leq C_p\|f\|_{p,(\alpha,\beta)} \tag{2.13}$$

implies

i) *the* $L^p(\gamma)$*-boundedness for the Gaussian-Riesz transform*

$$\|R^\gamma f\|_{p,\gamma} \leq C_p \|f\|_{p,\gamma}. \tag{2.14}$$

and

ii) *the* $L^p(\mu_\alpha)$*-boundedness for the Laguerre-Riesz transform*

$$\|R^\alpha f\|_{p,\alpha} \leq C_p \|f\|_{p,\alpha}. \tag{2.15}$$

Proof

i) Assume that the operator R^λ is bounded in $L^p\left([-1,1]\mu_\lambda\right)$. Let $\phi \in C_0^\infty(\mathbb{R})$, for each $\lambda > 0$ define the function

$$\phi_\lambda(x) = \phi(\sqrt{\lambda}x),$$

$x \in \mathbb{R}$. For λ big enough $sop\,\phi_\lambda$ is contained in $[-1,1]$. In what follows λ will be taken satisfying that condition.

Now, from the boundedness of R^λ we have

$$\|R^\lambda \phi_\lambda\|_{L^p([-1,1]\mu_\lambda)} \leq C\|\phi_\lambda\|_{L^p([-1,1]\mu_\lambda)},$$

in other words,

$$\left\| \sum_{n=1}^{\infty} \widehat{\phi_\lambda}(n) r_n^{(\lambda)} \sqrt{1-x^2}\, C_{n-1}^{\lambda+1} \right\|_{L^p([-1,1]\mu_\lambda)} \leq C\|\phi_\lambda\|_{L^p([-1,1]\mu_\lambda)},$$

where $r_n^{(\lambda)} = d_n^{(\lambda)} \nu_n^{-1/2} \frac{(n+2\lambda)}{2}$, $\nu_n = n(n+2\lambda)$ and $d_n^{(\lambda)} = \frac{4\Gamma(2\lambda)(n+\lambda)\Gamma(n+1)}{\Gamma(n+2\lambda+1)}$. Making the change of variable $x = \frac{y}{\sqrt{\lambda}}$ and taking $Z(\lambda) = \frac{\lambda^{1/2}[\Gamma(\lambda)]^2 2^{2\lambda}}{2\pi\Gamma(2\lambda)}$ we get

$$\left\{ \int_{-\sqrt{\lambda}}^{\sqrt{\lambda}} \left| \sum_{n=1}^{\infty} \widehat{\phi_\lambda}(n) r_n^{(\lambda)} \sqrt{1-\frac{y^2}{\lambda}}\, C_{n-1}^{\lambda+1}\left(\frac{y}{\sqrt{\lambda}}\right) \right|^p Z(\lambda)\left(1-\frac{y^2}{\lambda}\right)^{\lambda-1/2} dy \right\}^{1/p}$$
$$\leq C\|\phi_\lambda\|_{L^p([-1,1]\mu_\lambda)},$$

and therefore,

$$\left\{ \int_{-\sqrt{\lambda}}^{\sqrt{\lambda}} \left| \sum_{n=1}^{\infty} \widehat{\phi_\lambda}(n) r_n^{(\lambda)} \left(1-\frac{y^2}{\lambda}\right)^{\lambda/p-1/2p+1/2} e^{\frac{y^2}{p}} C_{n-1}^{\lambda+1}\left(\frac{y}{\sqrt{\lambda}}\right) \right|^p \frac{e^{-y^2}}{\sqrt{\pi}} dy \right\}^{1/p}$$
$$\leq C(Z(\lambda))^{-1/p}\|\phi_\lambda\|_{L^p([-1,1]\mu_\lambda)}.$$

On the other hand, we also have

$$\left\{ \int_{-\sqrt{\lambda}}^{\sqrt{\lambda}} \left| \sum_{n=1}^{\infty} \widehat{\phi_\lambda}(n) r_n^{(\lambda)} \left(1-\frac{y^2}{\lambda}\right)^{\lambda/2-1/4+1/2} e^{\frac{y^2}{2}} C_{n-1}^{\lambda+1}\left(\frac{y}{\sqrt{\lambda}}\right) \right|^2 \frac{e^{-y^2}}{\sqrt{\pi}} dy \right\}^{1/2}$$
$$\leq C(Z(\lambda))^{-1/2}\|\phi_\lambda\|_{L^2([-1,1]\mu_\lambda)}.$$

Define, for every $k \in \mathbb{N}$ and every $\lambda > 0$ such that $\sqrt{\lambda} > k$

$$F_{\lambda,k}(y) = \begin{cases} \sum_{n=1}^{\infty} \widehat{\phi_\lambda}(n) r_n^{(\lambda)} C_{n-1}^{\lambda+1}\left(\frac{y}{\sqrt{\lambda}}\right) \left(1-\frac{y^2}{\lambda}\right)^{\lambda/2-1/4+1/2} e^{\frac{y^2}{2}} & \text{if} \quad |y| \leq k \\ 0 & \text{if} \quad |y| > k, \end{cases}$$

and

$$f_{\lambda,k}(y) = \begin{cases} \sum\limits_{n=1}^{\infty} \widehat{\phi_\lambda}(n) r_n^{(\lambda)} C_{n-1}^{\lambda+1}(\frac{y}{\sqrt{\lambda}}) \left(1-\frac{y^2}{\lambda}\right)^{\lambda/p-1/2p+1/2} e^{\frac{y^2}{p}} & \text{if} \quad |y| \le k \\ 0 & \text{if} \quad |y| > k. \end{cases}$$

From the previous inequalities both series converges for all $y \in [-\sqrt{\lambda}, \sqrt{\lambda}]$, and $F_{\lambda,k} = f_{\lambda,k}\Omega_\lambda$, where

$$\Omega_\lambda(y) = e^{\frac{y^2}{2}-\frac{y^2}{p}} \left(1-\frac{y^2}{\lambda}\right)^{\lambda/2-\lambda/p-1/4+1/2p}$$

for all $k \in \mathbb{N}$ and $\sqrt{\lambda} > k$. Moreover

$$\begin{aligned} |\Omega_\lambda(y)| &= e^{\frac{y^2}{2}-\frac{y^2}{p}} \left(1-\frac{y^2}{\lambda}\right)^{\lambda/2-\lambda/p-1/4+1/2p} \\ &\le e^{\frac{y^2}{2}-\frac{y^2}{p}} e^{-\frac{y^2}{\lambda}(\lambda/2-\lambda/p-1/4+1/2p)} = e^{y^2(\frac{1}{4\lambda}-\frac{1}{2p\lambda})}. \end{aligned}$$

Therefore, if $p \le 2$ we have, $|\Omega_\lambda(y)| \le 1$ and for $p > 2$, $|\Omega_\lambda(y)| \le e^{k^2(\frac{1}{4\lambda}-\frac{1}{2p\lambda})}$. Thus Ω_λ is bounded in $[-k,k]$.
Now

$$(Z(\lambda))^{-1/p}\|\phi_\lambda\|_{L^p([-1,1]\mu_\lambda)} = (Z(\lambda))^{-1/p}\left\{\int_{-1}^{1} |\phi_\lambda(x)|^p \frac{\lambda\,[\Gamma(\lambda)]^2\, 2^{2\lambda}}{2\pi\Gamma\,(2\lambda)\,dx}\left(1-x^2\right)^{\lambda-1/2}\right\}^{1/p}$$

and therefore, by the change of variable $x = \frac{y}{\sqrt{\lambda}}$, we get

$$\begin{align} (Z(\lambda))^{-1/p}\|\phi_\lambda\|_{L^p([-1,1]\mu_\lambda)} &= (Z(\lambda))^{-1/p}\left\{\int_{-\sqrt{\lambda}}^{\sqrt{\lambda}} |\phi_\lambda(\frac{y}{\sqrt{\lambda}})|^p Z(\lambda)\left(1-\frac{y^2}{\lambda}\right)^{\lambda-1/2} dy\right\}^{1/p} \nonumber \\ &= \left\{\int_{-\sqrt{\lambda}}^{\sqrt{\lambda}} |\phi(y)|^p \left(1-\frac{y^2}{\lambda}\right)^{\lambda-1/2} dy\right\}^{1/p} \nonumber \\ &\le C\left\{\int_{-\sqrt{\lambda}}^{\sqrt{\lambda}} |\phi(y)|^p \frac{e^{-y^2}}{\sqrt{\pi}} dy\right\}^{1/p} \le C\|\phi\|_{L^p(\mathbb{R},\gamma)}, \tag{2.16} \end{align}$$

and then,

$$\begin{aligned} \lim_{\lambda\to\infty}(Z(\lambda))^{-1/p}\|\phi_\lambda\|_{L^p([-1,1]\mu_\lambda)} &\le \lim_{\lambda\to\infty} C\left\{\int_{-\sqrt{\lambda}}^{\sqrt{\lambda}} |\phi(y)|^p \frac{e^{-y^2}}{\sqrt{\pi}} dy\right\}^{1/p} \\ &= C\left\{\int_{-\infty}^{\infty} |\phi(y)|^p \frac{e^{-y^2}}{\sqrt{\pi}} dy\right\}^{1/p} = C\|\phi\|_{L^p(\mathbb{R},\gamma)}. \end{aligned}$$

On the other hand,

$$\begin{aligned} \|F_{\lambda,k}\|_{L^2(\mathbb{R},\gamma)} &= \left\{\int_{-k}^{k} |F_{\lambda,k}|^2 \frac{e^{-y^2}}{\sqrt{\pi}} dy\right\}^{1/2} \\ &\leq \left\{\int_{-\sqrt{\lambda}}^{\sqrt{\lambda}} |F_{\lambda,k}|^2 \frac{e^{-y^2}}{\sqrt{\pi}} dy\right\}^{1/2} \leq C(Z(\lambda))^{-1/2}\|\phi_\lambda\|_{L^2([-1,1]\mu_\lambda)}. \qquad (2.17) \end{aligned}$$

Thus, from (2.16) and (2.17), we have

$$\|F_{\lambda,k}\|_{L^2(\mathbb{R},\gamma)} \leq C\|\phi\|_{L^2(\mathbb{R},\gamma)}.$$

Similarly, as

$$\|F_{\lambda,k}\|_{L^p(\mathbb{R},\gamma)} = \left\{\int_{\mathbb{R}} |F_{\lambda,k}|^p \frac{e^{-y^2}}{\sqrt{\pi}} dy\right\}^{1/p} = \left\{\int_{-k}^{k} |f_{\lambda,k}\Omega_\lambda(y)|^p \frac{e^{-y^2}}{\sqrt{\pi}} dy\right\}^{1/p},$$

and using the boundedness of Ω_λ in $[-k,k]$, we get

$$\begin{aligned} \|F_{\lambda,k}\|_{L^p(\mathbb{R},\gamma)} &\leq C\left\{\int_{-k}^{k} |f_{\lambda,k}|^p \frac{e^{-y^2}}{\sqrt{\pi}} dy\right\}^{1/p} \\ &\leq C\left\{\int_{-\infty}^{\infty} |f_{\lambda,k}|^p \frac{e^{-y^2}}{\sqrt{\pi}} dy\right\}^{1/p} \leq C(Z(\lambda))^{-1/p}\|\phi_\lambda\|_{L^p([-1,1]\mu_\lambda)}. \qquad (2.18) \end{aligned}$$

Then, using (2.16) and (2.18) we get

$$\|F_{\lambda,k}\|_{L^p(\mathbb{R},\gamma)} \leq C\|\phi\|_{L^p(\mathbb{R},\gamma)}$$

for all $\sqrt{\lambda} > k$. Therefore $\{F_{\lambda,k}\}$ is a bounded sequence in $L^2(\mathbb{R},\gamma)$ and $L^p(\mathbb{R},\gamma)$, thus by the Bourbaki-Alaoglu's theorem, there exists an increasing sequence $\{\lambda_j\}_{j\in\mathbb{N}}$, such that $\lim_{j\to\infty}\lambda_j = \infty$, and functions $F_k \in L^2(\mathbb{R},\gamma)$ and $f_k \in L^p(\mathbb{R},\gamma)$ satisfying that

- $F_{\lambda,k} \to F_k$, as $j \to \infty$, in the weak topology on $L^2(\mathbb{R},\gamma)$
- $F_{\lambda,k} \to f_k$, as $j \to \infty$, in the weak topology on $L^p(\mathbb{R},\gamma)$

Moreover, $sopF_k \cup sopf_k \subseteq [-k,k]$, and

$$\|F_k\|_{L^2(\mathbb{R},\gamma)} \leq \lim_{j\to\infty} \|F_{\lambda_j,k}\|_{L^2(\mathbb{R},\gamma)} \leq C\|\phi\|_{L^2(\mathbb{R},\gamma)}, \qquad (2.19)$$

and

$$\|f_k\|_{L^p(\mathbb{R},\gamma)} \leq C\|\phi\|_{L^p(\mathbb{R},\gamma)}. \qquad (2.20)$$

Observe that, by Cauchy-Schwartz, defining for $k \in \mathbb{N}$,

$$\tau_k(g) = \int_{-\infty}^{\infty} g(x)\chi_{[-k,k]}(x)dx,$$

$\tau_k \in \left(L^2\left(\mathbb{R},\gamma\right)\right)^*$, and therefore,

$$\begin{aligned}\int_{-k}^{k} F_k(x)dx &= \int_{-\infty}^{\infty} F_k(x)\chi_{[-k,k]}(x)dx = \tau_k(F_k)\\ &= \lim_{j\to\infty} \tau_k(F_{\lambda_j,k}) = \lim_{j\to\infty} \int_{-\infty}^{\infty} F_{\lambda_j,k}(x)\chi_{[-k,k]}(x)dx\\ &= \int_{-\infty}^{\infty} f_k(x)\chi_{[-k,k]}(x)dx = \int_{-k}^{k} f_k(x)dx;\end{aligned}$$

i.e., $F_k = f_k \quad a.e \ (-k,k)$, for all k so $F_k = f_k \quad a.e \ \mathbb{R}$. Then from (2.20), we get

$$\|F_k\|_{L^p(\mathbb{R},\gamma)} \leq C\|\phi\|_{L^p(\mathbb{R},\gamma)}, \tag{2.21}$$

and therefore, from (2.19) and (2.21), there exists an increasing sequence $\{\lambda_j\}_{j\in\mathbb{N}} \subset (0,\infty)$, with $\lim_{j\to\infty}\lambda_j = \infty$,, and a function $F \in L^p\left(\mathbb{R},\gamma\right) \cap L^2\left(\mathbb{R},\gamma\right)$, such that

- For each $k \in \mathbb{N}$, $F_{\lambda_j,k} \to F$, as $j \to \infty$, in the weak topology of $L^2\left(\mathbb{R},\gamma\right)$ and in the weak topology of $L^p\left(\mathbb{R},\gamma\right)$, and
- $\|F\|_{L^p(\mathbb{R},\gamma)} \leq C\|\phi\|_{L^p(\mathbb{R},\gamma)}$.

For each $N \in \mathbb{N}$, $k \in \mathbb{N}$ and λ such that $\sqrt{\lambda} > k$, let us define

$$F_{\lambda,k}^N(y) = \chi_{[-k,k]}(y)\sum_{n=1}^{N}\widehat{\phi_\lambda}(n)r_n^{(\lambda)}C_{n-1}^{\lambda+1}(\frac{y}{\sqrt{\lambda}})\left(1-\frac{y^2}{\lambda}\right)^{\lambda/2-1/4+1/2} e^{\frac{y^2}{2}}$$

and

$$H_{\lambda,k}^N(y) = \chi_{[-k,k]}(y)\sum_{n=N+1}^{\infty}\widehat{\phi_\lambda}(n)r_n^{(\lambda)}C_{n-1}^{\lambda+1}(\frac{y}{\sqrt{\lambda}})\left(1-\frac{y^2}{\lambda}\right)^{\lambda/2-1/4+1/2} e^{\frac{y^2}{2}}.$$

Then, $F_{\lambda,k} = F_{\lambda,k}^N + H_{\lambda,k}^N$.

Now, we want to prove that for $k \in \mathbb{N}$ and $\lambda > 0$ such that $\sqrt{\lambda} > k$,

$$\int_{-\infty}^{\infty}\left|H_{\lambda,k}^N(y)\right|^2\frac{e^{-y^2}}{\sqrt{\pi}}dy \leq \frac{C}{N}, \tag{2.22}$$

uniformly in λ. Take $k \in \mathbb{N}$ and $\lambda > 0$; making the change of variable $x = \frac{y^2}{\lambda}$ and using Parseval's identity,

$$\begin{aligned}&\int_{-k}^{k}\left|H_{\lambda,k}^N(y)\right|^2\frac{e^{-y^2}}{\sqrt{\pi}}dy\\ &\quad\leq \int_{-\sqrt{\lambda}}^{\sqrt{\lambda}}\left|\sum_{n=N+1}^{\infty}\widehat{\phi_\lambda}(n)r_n^{(\lambda)}C_{n-1}^{\lambda+1}(\frac{y}{\sqrt{\lambda}})\left(1-\frac{y^2}{\lambda}\right)^{\lambda/2-1/4+1/2} e^{\frac{y^2}{2}}\right|^2\frac{e^{-y^2}}{\sqrt{\pi}}dy\\ &\quad= \frac{\sqrt{\lambda}}{\sqrt{\pi}}\int_{-1}^{1}\left|\sum_{n=N+1}^{\infty}\widehat{\phi_\lambda}(n)r_n^{(\lambda)}C_{n-1}^{\lambda+1}(x)\right|^2\left(1-x^2\right)^{\lambda+1/2}dx\\ &\quad= \frac{\sqrt{\lambda}}{\sqrt{\pi}}\frac{2\pi\Gamma(2\lambda+2)}{(\lambda+1)\left[\Gamma(\lambda+1)\right]^2 2^{2\lambda+2}}\sum_{n=N+1}^{\infty}\left|\widehat{\phi_\lambda}(n)\right|^2\left|r_n^{(\lambda)}\right|^2\left\|C_{n-1}^{\lambda+1}\right\|_{2,(\lambda+1)}^2.\end{aligned}$$

Using (1.16) we have

$$-\frac{2\lambda}{n(n+2\lambda)}\frac{d}{dx}\left\{(1-x^2)^{\lambda+1/2}C_{n-1}^{\lambda+1}(x)\right\} = C_n^{\lambda}(x)\,(1-x^2)^{\lambda-1/2}.$$

Now, integrating by parts, we have

$$\begin{aligned}
\widehat{\phi_\lambda}(n) &= \frac{\lambda\,[\Gamma(\lambda)]^2\,2^{2\lambda}}{2\pi\Gamma(2\lambda)}\left\{\frac{\sqrt{\lambda}2\lambda}{n(n+2\lambda)}\int_{-1}^{1}\phi'(\sqrt{\lambda}x)C_{n-1}^{\lambda+1}(x)(1-x^2)^{\lambda+1/2}dx\right\}\\
&= \frac{(2\lambda+1)}{(\lambda+1)}\frac{\lambda^{3/2}}{n(n+2\lambda)}\\
&\qquad\times\left\{\frac{(\lambda+1)\,[\Gamma(\lambda+1)]^2\,2^{2\lambda+2}}{2\pi\Gamma(2\lambda+2)}\int_{-1}^{1}\phi'(\sqrt{\lambda}x)C_{n-1}^{\lambda+1}(x)(1-x^2)^{\lambda+1/2}dx\right\}\\
&= \frac{(2\lambda+1)}{(\lambda+1)}\frac{\lambda^{3/2}}{n(n+2\lambda)}\left\langle\phi'(\sqrt{\lambda}\cdot),C_{n-1}^{\lambda+1}\right\rangle_{(\lambda+1)}.
\end{aligned}$$

Then,

$$\begin{aligned}
&\int_{-k}^{k}\left|H_{\lambda,k}^{N}(y)\right|^2\frac{e^{-y^2}}{\sqrt{\pi}}dy\\
&\quad\le\frac{\sqrt{\lambda}}{\sqrt{\pi}}\frac{2\pi\Gamma(2\lambda+2)}{(\lambda+1)\,[\Gamma(\lambda+1)]^2\,2^{2\lambda+2}}\\
&\qquad\times\sum_{n=N+1}^{\infty}\frac{(2\lambda+1)^2}{(\lambda+1)^2}\frac{\lambda^3}{n^2(n+2\lambda)^2}\left|\left\langle\phi'(\sqrt{\lambda}\cdot),C_{n-1}^{\lambda+1}\right\rangle_{(\lambda+1)}\right|^2\left|r_n^{(\lambda)}\right|^2\left\|C_{n-1}^{\lambda+1}\right\|_{2,(\lambda+1)}^2\\
&\quad\le\frac{\sqrt{\lambda}}{\sqrt{\pi}}\frac{2\pi\Gamma(2\lambda+2)}{(\lambda+1)\,[\Gamma(\lambda+1)]^2\,2^{2\lambda+2}}\\
&\qquad\times\sum_{n=N+1}^{\infty}\frac{(2\lambda+1)^2}{(\lambda+1)^2}\frac{\lambda^3}{n^2(n+2\lambda)^2}\left|\frac{\left\langle\phi'(\sqrt{\lambda}\cdot),C_{n-1}^{\lambda+1}\right\rangle_{(\lambda+1)}}{\left\|C_{n-1}^{\lambda+1}\right\|_{2,(\lambda+1)}}\right|^2\left|r_n^{(\lambda)}\right|^2\left\|C_{n-1}^{\lambda+1}\right\|_{2,(\lambda+1)}^4.
\end{aligned}$$

Now, using (1.15) and that $r_n^{(\lambda)}=\frac{2\Gamma(2\lambda)(n+\lambda)\Gamma(n+1)}{\Gamma(n+2\lambda+1)}\left(\frac{(n+2\lambda)}{n}\right)^{1/2}$, we get

$$\begin{aligned}
\left|r_n^{(\lambda)}\right|^2\left\|C_{n-1}^{\lambda+1}\right\|_{2,(\lambda+1)}^4 &= \frac{4\,[\Gamma(2\lambda)]^2\,(n+\lambda)^2(n!)^2}{[\Gamma(n+2\lambda+1)]^2}\frac{(n+2\lambda)}{n}\\
&\qquad\times\frac{(\lambda+1)^2\,[\Gamma(n+2\lambda+1)]^2\,n^2}{(2\lambda+1)^2\,[\Gamma(2\lambda+1)]^2\,(n+\lambda)^2(n!)^2}\\
&= \frac{4\,[\Gamma(2\lambda)]^2\,(n+2\lambda)(\lambda+1)^2n}{(2\lambda+1)^2\,[\Gamma(2\lambda+1)]^2}=\frac{(\lambda+1)^2(n+2\lambda)n}{(2\lambda+1)^2\lambda^2}.
\end{aligned}$$

Therefore,

$$\begin{aligned}
&\int_{-k}^{k}\left|H_{\lambda,k}^{N}(y)\right|^{2}\frac{e^{-y^{2}}}{\sqrt{\pi}}dy\\
&\quad\leq \frac{\sqrt{\lambda}}{\sqrt{\pi}}\frac{2\pi\Gamma(2\lambda+2)}{(\lambda+1)\left[\Gamma(\lambda+1)\right]^{2}2^{2\lambda+2}}\sum_{n=N+1}^{\infty}\frac{\lambda}{n(n+2\lambda)}\left|\frac{\left\langle\phi'(\sqrt{\lambda}\cdot),C_{n-1}^{\lambda+1}\right\rangle_{(\lambda+1)}}{\left\|C_{n-1}^{\lambda+1}\right\|_{2,(\lambda+1)}}\right|^{2}\\
&\quad\leq \frac{\sqrt{\lambda}}{\sqrt{\pi}}\frac{2\pi\Gamma(2\lambda+2)}{(\lambda+1)\left[\Gamma(\lambda+1)\right]^{2}2^{2\lambda+2}}\frac{1}{N}\sum_{n=1}^{\infty}\left|\frac{\left\langle\phi'(\sqrt{\lambda}\cdot),C_{n-1}^{\lambda+1}\right\rangle_{(\lambda+1)}}{\left\|C_{n-1}^{\lambda+1}\right\|_{2,(\lambda+1)}}\right|^{2},
\end{aligned}$$

but,

$$\begin{aligned}
&\sum_{n=1}^{\infty}\left|\frac{\left\langle\phi'(\sqrt{\lambda}\cdot),C_{n-1}^{\lambda+1}\right\rangle_{(\lambda+1)}}{\left\|C_{n-1}^{\lambda+1}\right\|_{2,(\lambda+1)}}\right|^{2}\\
&\quad=\frac{(\lambda+1)\left[\Gamma(\lambda+1)\right]^{2}2^{2\lambda+2}}{2\pi\Gamma(2\lambda+2)}\int_{-1}^{1}\left|\phi'(\sqrt{\lambda}x)\right|^{2}(1-x^{2})^{\lambda+1/2}dx\\
&\quad=\frac{(\lambda+1)\left[\Gamma(\lambda+1)\right]^{2}2^{2\lambda+2}\lambda^{-1/2}}{2\pi\Gamma(2\lambda+2)}\int_{-\sqrt{\lambda}}^{\sqrt{\lambda}}\left|\phi'(y)\right|^{2}(1-\frac{y^{2}}{\sqrt{\lambda}})^{\lambda+1/2}dy\\
&\quad\leq C\frac{\sqrt{\pi}(\lambda+1)\left[\Gamma(\lambda+1)\right]^{2}2^{2\lambda+2}\lambda^{-1/2}}{2\pi\Gamma(2\lambda+2)}\int_{-\infty}^{\infty}\left|\phi'(y)\right|^{2}\frac{e^{-y^{2}}}{\sqrt{\pi}}dy
\end{aligned}$$

Then,

$$\int_{-k}^{k}\left|H_{\lambda,k}^{N}(y)\right|^{2}\frac{e^{-y^{2}}}{\sqrt{\pi}}dy\leq\frac{C}{N}\left\|\phi'\right\|_{L^{2}(\mathbb{R},\gamma)}^{2}$$

i.e.,

$$\int_{-\infty}^{\infty}\left|H_{\lambda,k}^{N}(y)\right|^{2}\frac{e^{-y^{2}}}{\sqrt{\pi}}dy \;=\; \int_{-k}^{k}\left|H_{\lambda,k}^{N}(y)\right|^{2}\frac{e^{-y^{2}}}{\sqrt{\pi}}dy\leq\frac{C}{N}.$$

Thus $\{H_{\lambda,k}^{N}\}_{j\in\mathbb{N}}$ is a bounded sequence on $L^{2}(\mathbb{R},\gamma)$, so by Bourbaki-Alaoglu's theorem, there exists a sequence $(\lambda_j)_{j\in\mathbb{N}}$, $\lim_{j\to\infty}\lambda_j=\infty$ such that, for all $N\in\mathbb{N}$, $\{H_{\lambda_j,k}^{N}\}_{j\in\mathbb{N}}$ converges weakly in $L^{2}(\mathbb{R},\gamma)$ to a function $H_{k}^{N}\in L^{2}(\mathbb{R},\gamma)$. Moreover,

$$\int_{-\infty}^{\infty}\left|H_{k}^{N}(y)\right|^{2}\frac{e^{-y^{2}}}{\sqrt{\pi}}dy\leq\frac{C}{N}. \tag{2.23}$$

Then, there exists a non decreasing sequence $(N_j)_{j\in\mathbb{N}}$ such that,

$$H_{k}^{N_j}(y)\longrightarrow 0,\ \ a.e.\ \ y\in\mathbb{R}. \tag{2.24}$$

Defining, for each $m\in\mathbb{N}$, $F_{k}^{N_m}=F-H_{k}^{N_m}$; then since $F_{\lambda,k}\to F_k$, as $j\to\infty$ in the weak topology of $L^{2}(\mathbb{R},\gamma)$ and that $F_{\lambda,k}^{N}=F_{\lambda,k}-H_{\lambda,k}^{N}$, we have, for all $m \in \mathbb{N}$,

$$F^{N_m}_{\lambda_j,k} \longrightarrow F^{N_m}_{k}, \qquad \text{weakly in } \; L^2(\mathbb{R},\gamma) \quad \text{as} \quad j \to \infty. \tag{2.25}$$

Also, from (2.24) we get that

$$F^{N_m}_{k}(y) \longrightarrow F(y), \;\; \text{as } m \to \infty, \;\; a.e. \;\; y \in \mathbb{R}. \tag{2.26}$$

To finish the proof of i) we need to prove that F is the Hermite-Riesz transform of ϕ, i.e.

$$F(y) = \sum_{n=1}^{\infty} \widehat{\phi}^{\gamma}(n) \frac{1}{2^n n!} \sqrt{2n} H_{n-1}(y), \tag{2.27}$$

where

$$\widehat{\phi}^{\gamma}(n) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} \phi(y) H_n(y) e^{-y^2} dy,$$

is the n-th Fourier-Hermite coefficient of ϕ. Observe that,

$$\begin{aligned} F^{N}_{\lambda,k}(y) &= \sum_{n=1}^{N} \widehat{\phi_{\lambda}}(n) \frac{2\Gamma(2\lambda)(n+\lambda)n!(n+2\lambda)^{1/2}}{\Gamma(n+2\lambda+1)n^{1/2}} C^{\lambda+1}_{n-1}(\frac{y}{\sqrt{\lambda}}) \left(1 - \frac{y^2}{\lambda}\right)^{\lambda/2+1/4} e^{\frac{y^2}{2}} \\ &= \sum_{n=1}^{N} \lambda^{-n/2} \widehat{\phi_{\lambda}}(n) \frac{(\frac{n}{\lambda}+1)n!}{n^{1/2}(\frac{n}{\lambda}+2)^{1/2} 2(2+\frac{1}{\lambda})(2+\frac{2}{\lambda})\dots(2+\frac{n-1}{\lambda})} \\ &\qquad \times 2\lambda^{-\frac{(n-1)}{2}} C^{\lambda+1}_{n-1}(\frac{y}{\sqrt{\lambda}}) \left(1 - \frac{y^2}{\lambda}\right)^{\lambda/2+1/4} e^{\frac{y^2}{2}}. \end{aligned} \tag{2.28}$$

From the asymptotic relation between Jacobi and Hermite polynomials (1.33), it is easy to see that

$$\begin{aligned} \lim_{\lambda\to\infty} \lambda^{-\frac{(n-1)}{2}} C^{\lambda+1}_{n-1}\left(\frac{y}{\sqrt{\lambda}}\right) &= \lim_{\lambda\to\infty} (\lambda+1)^{-\frac{(n-1)}{2}} \left(1+\frac{1}{\lambda}\right)^{\frac{(n-1)}{2}} C^{\lambda+1}_{n-1}\left(\frac{y}{\sqrt{\lambda+1}}\sqrt{1+\frac{1}{\lambda}}\right) \\ &= \frac{H_{n-1}(y)}{(n-1)!}, \end{aligned}$$

uniformly in $[-k,k]$. On the other hand, observe

$$\lim_{\lambda\to\infty} \lambda^{-n/2} \widehat{\phi_{\lambda}}(n) = \frac{1}{\sqrt{\pi} n!} \int_{-\infty}^{\infty} \phi(y) H_n(y) e^{-y^2} dy = \frac{1}{n!} \widehat{\phi}^{\gamma}(n).$$

Then taking limit as $\lambda \to \infty$ in (2.28) we get

$$\begin{aligned} \lim_{\lambda\to\infty} F^{N}_{\lambda,k}(y) &= \sum_{n=1}^{N} \frac{1}{n!} \widehat{\phi}^{\gamma}(n) \frac{n!}{n^{1/2} 2^{1/2} 2^n} 2 \frac{H_{n-1}(y)}{(n-1)!} \\ &= \sum_{n=1}^{N} \widehat{\phi}^{\gamma}(n) \frac{1}{2^n n!} \sqrt{2n} H_{n-1}(y). \end{aligned}$$

Hence,

$$F^{N_m}_{k}(y) = \sum_{n=1}^{N_m} \widehat{\phi}^{\gamma}(n) \frac{1}{2^n n!} \sqrt{2n} H_{n-1}(y), \tag{2.29}$$

and from (2.26) we finally get (2.27),

$$F(y)=\sum_{n=1}^{\infty}\widehat{\phi}^{\gamma}(n)\frac{1}{2^{n}n!}\sqrt{2n}H_{n-1}(y).$$

ii) The proof of ii) is essentially analogous to i), so we will give less details in this case. Assume that the operator $R^{\alpha,\beta}$ is bounded in $L^p\left([-1,1]\mu_{\alpha,\beta}\right)$. Let $\phi\in C_0^{\infty}((0,\infty))$ and $\beta>0$, then we have

$$\|R^{\alpha,\beta}\phi_\beta\|_{L^p\left([-1,1]\mu_{(\alpha,\beta)}\right)}\leq C\|\phi_\beta\|_{L^p\left([-1,1]\mu_{(\alpha,\beta)}\right)},$$

where $\phi_\beta(x)=\phi\left(\frac{\beta}{2}(1-x)\right)$, for $x\in(0,\infty)$, i.e.

$$\left\{\eta_{\alpha,\beta}\int_{-1}^{1}\left|\sum_{n=1}^{\infty}\frac{\widehat{\phi_\beta}(n)}{\hat{h}_n^{(\alpha,\beta)}}\frac{(n+\alpha+\beta+1)^{1/2}}{2n^{1/2}}\sqrt{1-x^2}P_{n-1}^{(\alpha+1,\beta+1)}(x)\right|^p(1-x)^{\alpha}(1+x)^{\beta}dx\right\}^{1/p}$$
$$\leq C\|\phi_\beta\|_{L^p\left([-1,1]\mu_{(\alpha,\beta)}\right)}.$$

Now, making the change of variable $x=1-\frac{2}{\beta}y$

$$\left\{\eta_{\alpha,\beta}\frac{2^{\alpha+\beta+1}}{\beta^{\alpha+1}}\int_0^{\beta}\left|\sum_{n=1}^{\infty}\frac{\widehat{\phi_\beta}(n)}{\hat{h}_n^{(\alpha,\beta)}}\frac{(n+\alpha+\beta+1)^{1/2}}{n^{1/2}}P_{n-1}^{(\alpha+1,\beta+1)}\left(1-\frac{2}{\beta}y\right)\left(\frac{y}{\beta}\right)^{1/2}\right.\right.$$
$$\left.\left.\times\left(1-\frac{y}{\beta}\right)^{1/2+\beta/p}e^{\frac{y}{p}}\right|^p y^{\alpha}e^{-y}dx\right\}^{1/p}\leq C\|\phi_\beta\|_{L^p\left([-1,1]\mu_{(\alpha,\beta)}\right)},$$

and therefore

$$\left\{\int_0^{\beta}\left|\sum_{n=1}^{\infty}\frac{\widehat{\phi_\beta}(n)}{\hat{h}_n^{(\alpha,\beta)}}\frac{(n+\alpha+\beta+1)^{1/2}}{n^{1/2}}P_{n-1}^{(\alpha+1,\beta+1)}\left(1-\frac{2}{\beta}y\right)\left(\frac{y}{\beta}\right)^{1/2}\right.\right.$$
$$\left.\left.\times\left(1-\frac{y}{\beta}\right)^{1/2+\beta/p}e^{\frac{y}{p}}\right|^p y^{\alpha}e^{-y}dx\right\}^{1/p}\leq C(Z_{\alpha,\beta})^{-1/p}\|\phi_\beta\|_{L^p\left([-1,1]\mu_{(\alpha,\beta)}\right)}$$

where $Z_{\alpha,\beta}=\frac{\Gamma(\alpha+\beta+2)}{\Gamma(\alpha+1)\beta^{\alpha+2}\Gamma(\beta)}$. Analogously we get,

$$\left\{\int_0^{\beta}\left|\sum_{n=1}^{\infty}\frac{\widehat{\phi_\beta}(n)}{\hat{h}_n^{(\alpha,\beta)}}\frac{(n+\alpha+\beta+1)^{1/2}}{n^{1/2}}P_{n-1}^{(\alpha+1,\beta+1)}\left(1-\frac{2}{\beta}y\right)\left(\frac{y}{\beta}\right)^{1/2}\right.\right.$$
$$\left.\left.\times\left(1-\frac{y}{\beta}\right)^{1/2+\beta/2}e^{\frac{y}{2}}\right|^2 y^{\alpha}e^{-y}dx\right\}^{1/2}\leq C(Z_{\alpha,\beta})^{-1/2}\|\phi_\beta\|_{L^2\left([-1,1]\mu_{(\alpha,\beta)}\right)}.$$

Now, for each $k\in\mathbb{N}$ and $\beta>0$, such that $\beta>k$, define the functions

$$F_{\beta,k}(y) = -\chi_{(0,k)}(y)\sum_{n=1}^{\infty}\frac{\widehat{\phi_\beta}(n)}{\hat{h}_n^{(\alpha,\beta)}}\frac{(n+\alpha+\beta+1)^{1/2}}{n^{1/2}}P_{n-1}^{(\alpha+1,\beta+1)}\left(1-\frac{2}{\beta}y\right)$$
$$\times\left(\frac{y}{\beta}\right)^{1/2}\left(1-\frac{y}{\beta}\right)^{1/2+\beta/2}e^{\frac{y}{2}},$$

and

$$\begin{aligned} f_{\beta,k}(y) &= -\chi_{(0,k)}(y)\sum_{n=1}^{\infty}\frac{\widehat{\phi_\beta}(n)}{\hat{h_n}^{(\alpha,\beta)}}\frac{(n+\alpha+\beta+1)^{1/2}}{n^{1/2}}P_{n-1}^{(\alpha+1,\beta+1)}\left(1-\frac{2}{\beta}y\right)\\ &\qquad\times\left(\frac{y}{\beta}\right)^{1/2}\left(1-\frac{y}{\beta}\right)^{1/2+\beta/p}e^{\frac{y}{p}}. \end{aligned}$$

From the previous inequalities both series converge for all $y \in (0,\beta)$, and $F_{\beta,k} = f_{\beta,k}\Omega_\beta$, where

$$\Omega_\beta(y) = e^{\frac{y}{2}-\frac{y}{p}}\left(1-\frac{y}{\beta}\right)^{\beta/2-\beta/p}$$

for all $k \in \mathbb{N}$ and $\beta > 0$. Now as,

$$|\Omega_\beta(y)| = e^{\frac{y}{2}-\frac{y}{p}}\left(1-\frac{y}{\beta}\right)^{\beta/2-\beta/p} \leq e^{\frac{y}{2}-\frac{y}{p}}(e^{-y/\beta})^{\beta/2-\beta/p} = 1$$

we conclude that Ω_β is bounded in $(0,k)$. On the other hand,

$$(Z_{\alpha,\beta})^{-1/p}\|\phi_\beta\|_{L^p\left([-1,1]\mu_{(\alpha,\beta)}\right)} = (Z_{\alpha,\beta})^{-1/p}\left\{\eta_{\alpha,\beta}\int_{-1}^{1}\left|\widehat{\phi_\beta}(x)\right|^p(1-x)^\alpha(1+x)^\beta dx\right\}^{1/p},$$

and making the change of variable $x = 1-\frac{2}{\beta}y$ we get

$$\begin{aligned} &(Z_{\alpha,\beta})^{-1/p}\left\{\eta_{\alpha,\beta}\int_{-1}^{1}|\phi_\beta(x)|^p(1-x)^\alpha(1+x)^\beta dx\right\}^{1/p}\\ &= (Z_{\alpha,\beta})^{-1/p}\left\{\eta_{\alpha,\beta}\frac{2}{\beta}\int_0^\beta\left|\phi_\beta(1-\frac{2}{\beta}y)\right|^p\left(\frac{2y}{\beta}\right)^\alpha\left(2-\frac{2y}{\beta}\right)^\beta dy\right\}^{1/p}\\ &= \left\{\int_0^\beta|\phi(y)|^p y^\alpha\left(1-\frac{y}{\beta}\right)^\beta dy\right\}^{1/p} \leq \left\{\int_0^\infty|\phi(y)|^p y^\alpha e^{-y}dy\right\}^{1/p} = \|\phi\|_{L^p((0,\infty),\mu_\alpha)}. \end{aligned}$$

Then,

$$\begin{aligned} \lim_{\beta\to\infty} C(Z_{\alpha,\beta})^{-1/p}\|\phi_\beta\|_{L^p\left([-1,1]\mu_{(\alpha,\beta)}\right)} &\leq \lim_{\beta\to\infty} C\left\{\int_0^\infty|\phi(y)|^p y^\alpha e^{-y}dy\right\}^{1/p}\\ &= C\|\phi\|_{L^p((0,\infty),\mu_\alpha)}. \end{aligned}$$

Moreover,

$$(Z_{\alpha,\beta})^{-1/p}\|\phi_\beta\|_{L^p\left([-1,1],\mu_{(\alpha,\beta)}\right)} \leq C\|\phi\|_{L^p((0,\infty),\mu_\alpha)}. \tag{2.30}$$

Now,

$$\begin{aligned} \|F_{\beta,k}\|_{L^2((0,\infty),\mu_\alpha)} &= \left\{\frac{1}{\Gamma(\alpha+1)}\int_0^k|F_{\beta,k}|^2y^\alpha e^{-y}dy\right\}^{1/2}\\ &\leq C(Z_{\alpha,\beta})^{-1/2}\|\phi_\beta\|_{L^2\left([-1,1]\mu_{(\alpha,\beta)}\right)} \end{aligned}$$

and therefore, from (2.30) for $p = 2$ and (2.31), we get

$$\|F_{\beta,k}\|_{L^2((0,\infty),\mu_\alpha)} \leq C\|\phi\|_{L^2((0,\infty),\mu_\alpha)}.$$

Analogously, using that Ω_β is bounded in $(0,k)$,

$$\begin{aligned}\|F_{\beta,k}\|_{L^p((0,\infty),\mu_\alpha)} &= \left\{\frac{1}{\Gamma(\alpha+1)}\int_0^k |f_{\beta,k}\Omega_\beta(y)|^p y^\alpha e^{-y}dy\right\}^{1/p}\\ &\leq C\left\{\frac{1}{\Gamma(\alpha+1)}\int_0^k |f_{\beta,k}|^p y^\alpha e^{-y}dy\right\}^{1/p}\\ &\leq C(Z_{\alpha,\beta})^{-1/p}\|\phi_\beta\|_{L^p\left([-1,1]\mu_{(\alpha,\beta)}\right)}. \qquad (2.31)\end{aligned}$$

Then, from (2.30) and (2.31),

$$\|F_{\beta,k}\|_{L^p((0,\infty),\mu_\alpha)} \leq C\|\phi\|_{L^p((0,\infty),\mu_\alpha)},$$

for all $\beta > k$. Therefore $\{F_{\beta,k}\}$ is a bounded subsequence in $L^2\left((0,\infty),\mu_\alpha\right)$ with $L^p\left((0,\infty),\mu_\alpha\right)$. Thus by the Bourbaki-Alaoglu's theorem, there exists an increasing sequence $\{\beta_j\}_{j\in\mathbb{N}}$ with $\lim_{j\to\infty}\beta_j = \infty$ and functions $F_k \in L^2\left((0,\infty),\mu_\alpha\right)$ and $f_k \in L^p\left((0,\infty),\mu_\alpha\right)$ satisfying that

- $F_{\beta,k} \to F_k$, as $j\to\infty$, in the weak topology of $L^2\left((0,\infty),\mu_\alpha\right)$
- $F_{\beta,k} \to f_k$, as $j\to\infty$, in the weak topology of $L^p\left((0,\infty),\mu_\alpha\right)$.

Then, as in i), we can conclude that there exists an increasing sequence $(\beta_j)_{j\in\mathbb{N}} \subset (0,\infty)$ such that $\lim_{j\to\infty}\beta_j = \infty$, and a function $F \in L^p\left((0,\infty),\mu_\alpha\right)\cap L^2\left((0,\infty),\mu_\alpha\right)$, such that

- for each $k\in\mathbb{N}$, $F_{\beta_j,k}\to F$, as $j\to\infty$, in the weak topology on $L^2\left((0,\infty),\mu_\alpha\right)$ and in the weak topology on $L^p\left((0,\infty),\mu_\alpha\right)$.
- $\|F\|_{L^p((0,\infty),\mu_\alpha)} \leq C\|\phi\|_{L^p((0,\infty),\mu_\alpha)}$.

For each $N\in\mathbb{N}$, $k\in\mathbb{N}$ and β such that $\beta > k$, let us define

$$\begin{aligned}F^N_{\beta,k}(y) = -\chi_{(0,k)}(y)\sum_{n=1}^{N}\frac{\widehat{\phi_\beta}(n)}{\hat{h}_n^{(\alpha,\beta)}}\frac{(n+\alpha+\beta+1)^{1/2}}{n^{1/2}}P^{(\alpha+1,\beta+1)}_{n-1}\left(1-\frac{2}{\beta}y\right)\\ \times\left(\frac{y}{\beta}\right)^{1/2}\left(1-\frac{y}{\beta}\right)^{1/2+\beta/2}e^{\frac{y}{2}}\end{aligned}$$

and

$$\begin{aligned}H^N_{\beta,k}(y) = -\chi_{(0,k)}(y)\sum_{n=N+1}^{\infty}\frac{\widehat{\phi_\beta}(n)}{\hat{h}_n^{(\alpha,\beta)}}\frac{(n+\alpha+\beta+1)^{1/2}}{n^{1/2}}P^{(\alpha+1,\beta+1)}_{n-1}\left(1-\frac{2}{\beta}y\right)\\ \times\left(\frac{y}{\beta}\right)^{1/2}\left(1-\frac{y}{\beta}\right)^{1/2+\beta/2}e^{\frac{y}{2}}.\end{aligned}$$

Then, $F_{\beta,k} = F^N_{\beta,k} + H^N_{\beta,k}$.
Again, we want to prove that for $k\in\mathbb{N}$ and $\lambda > 0$ such that $\sqrt{\lambda} > k$,

$$\int_0^\infty \left|H^N_{\beta,k}(y)\right|^2 y^\alpha e^{-y}dy \leq \frac{C}{N}. \qquad (2.32)$$

uniformly in β. Take $k \in \mathbb{N}$, and β such that $\beta > k$, then using Parseval's identity, the change of variable $x = 1 - \frac{2}{\beta}y$ and (1.4),

$$\begin{aligned}
&\frac{1}{\Gamma(\alpha+1)}\int_0^\infty |H_{\beta,k}^N(y)|^2 y^\alpha e^{-y}dy \\
&= \frac{\beta^\alpha}{\Gamma(\alpha+1)}\int_0^\beta \left|\sum_{n=N+1}^\infty \frac{\widehat{\phi_\beta}(n)}{\hat{h}_n^{(\alpha,\beta)}}\frac{(n+\alpha+\beta+1)^{1/2}}{n^{1/2}}P_{n-1}^{(\alpha+1,\beta+1)}\left(1-\frac{2}{\beta}y\right)\right|^2 \\
&\qquad\qquad\qquad\qquad \times\left(\frac{y}{\beta}\right)^{\alpha+1}\left(1-\frac{y}{\beta}\right)^{\beta+1}dy \\
&= \frac{\beta^{\alpha+1}}{2^{\alpha+\beta+3}\Gamma(\alpha+1)}\int_{-1}^1 \left|\sum_{n=N+1}^\infty \frac{\widehat{\phi_\beta}(n)}{\hat{h}_n^{(\alpha,\beta)}}\frac{(n+\alpha+\beta+1)^{1/2}}{n^{1/2}}P_{n-1}^{(\alpha+1,\beta+1)}(x)\right|^2 \\
&\qquad\qquad\qquad\qquad \times(1-x)^{\alpha+1}(1+x)^{\beta+1}dx \\
&= \frac{\beta^{\alpha+1}B(\alpha+2,\beta+2)}{\Gamma(\alpha+1)}\sum_{n=N+1}^\infty\left|\frac{\widehat{\phi_\beta}(n)}{\hat{h}_n^{(\alpha,\beta)}}\right|^2\frac{(n+\alpha+\beta+1)}{n}\left\|P_{n-1}^{(\alpha+1,\beta+1)}\right\|^2_{\mu_{(\alpha+1,\beta+1)}} \\
&= \frac{\beta^{\alpha+1}B(\alpha+2,\beta+2)}{\Gamma(\alpha+1)}\sum_{n=N+1}^\infty\left|\frac{\widehat{\phi_\beta}(n)}{\hat{h}_n^{(\alpha,\beta)}}\right|^2\frac{(n+\alpha+\beta+1)}{n} \\
&\qquad\qquad \times\frac{(\alpha+\beta+3)(\alpha+\beta+2)n}{(\alpha+1)(\beta+1)\,(n+\alpha+\beta+1)}\left\|P_n^{(\alpha,\beta)}\right\|^2_{\mu_{(\alpha,\beta)}} \\
&= \frac{\beta^{\alpha+1}B(\alpha+2,\beta+2)}{\Gamma(\alpha+1)}\sum_{n=N+1}^\infty\frac{\left|\widehat{\phi_\beta}(n)\right|^2}{\hat{h}_n^{(\alpha,\beta)}}\frac{(\alpha+\beta+3)(\alpha+\beta+2)}{(\alpha+1)(\beta+1)}.
\end{aligned}$$

Now, integrating by parts and using (1.5)

$$\begin{aligned}
\widehat{\phi_\beta}(n) &= -\frac{\beta}{2^{\alpha+\beta+3}B(\alpha+1,\beta+1)n}\int_{-1}^1 \phi'\left(\frac{\beta}{2}(1-x)\right)P_{n-1}^{(\alpha+1,\beta+1)}(x) \\
&\qquad\qquad\qquad\qquad \times(1-x)^{\alpha+1}(1+x)^{\beta+1}dx \\
&= -\frac{\beta(\alpha+1)(\beta+1)}{2^{\alpha+\beta+3}(\alpha+\beta+3)(\alpha+\beta+2)B(\alpha+2,\beta+2)n}\int_{-1}^1\phi'\left(\frac{\beta}{2}(1-x)\right) \\
&\qquad\qquad\qquad\qquad \times P_{n-1}^{(\alpha+1,\beta+1)}(x)\,(1-x)^{\alpha+1}(1+x)^{\beta+1}dx \\
&= -\frac{\beta(\alpha+1)(\beta+1)}{n(\alpha+\beta+3)(\alpha+\beta+2)}\left\langle\phi'\left(\frac{\beta}{2}(1-\cdot)\right), P_{n-1}^{(\alpha+1,\beta+1)}\right\rangle_{(\alpha+1,\beta+1)}.
\end{aligned}$$

Then,

$$\frac{1}{\Gamma(\alpha+1)}\int_0^k |H_{\beta,k}^N(y)|^2 y^\alpha e^{-y}dy$$

$$
\begin{aligned}
&= \frac{\beta^{\alpha+3} B(\alpha+2,\beta+2)}{\Gamma(\alpha+1)} \\
&\qquad \times \sum_{n=N+1}^{\infty} \frac{\left|\left\langle \phi'\left(\frac{\beta}{2}(1-\cdot)\right), P_{n-1}^{(\alpha+1,\beta+1)}\right\rangle_{(\alpha+1,\beta+1)}\right|^2}{\left\|P_{n-1}^{(\alpha+1,\beta+1)}\right\|^2_{\mu_{(\alpha+1,\beta+1)}}} \frac{1}{n(n+\alpha+\beta+1)} \\
&\le \frac{\beta^{\alpha+2} B(\alpha+2,\beta+2)}{\Gamma(\alpha+1)N} \sum_{n=N+1}^{\infty} \frac{\left|\left\langle \phi'\left(\frac{\beta}{2}(1-\cdot)\right), P_{n-1}^{(\alpha+1,\beta+1)}\right\rangle_{(\alpha+1,\beta+1)}\right|^2}{\left\|P_{n-1}^{(\alpha+1,\beta+1)}\right\|^2_{\mu_{(\alpha+1,\beta+1)}}}.
\end{aligned}
$$

But, making the change of variable $y = \frac{\beta}{2}(1-x)$, i.e. $x = 1 - \frac{2y}{\beta}$,

$$
\begin{aligned}
&\sum_{n=N+1}^{\infty} \frac{\left|\left\langle \phi'\left(\frac{\beta}{2}(1-\cdot)\right), P_{n-1}^{(\alpha+1,\beta+1)}\right\rangle_{(\alpha+1,\beta+1)}\right|^2}{\left\|P_{n-1}^{(\alpha+1,\beta+1)}\right\|^2_{\mu_{(\alpha+1,\beta+1)}}} \\
&= \frac{1}{B(\alpha+2,\beta+2)} \frac{\beta}{\beta^{\alpha+2}} \int_0^{\beta} \left|\phi'(y)\right|^2 y^{\alpha+1} \left(1-\frac{y}{\beta}\right)^{\beta+1} dy \\
&\le \frac{1}{B(\alpha+2,\beta+2)} \frac{1}{\beta^{\alpha+2}} \int_0^{\infty} \left|\phi'(y)\right|^2 y^{\alpha+1} e^{-y} dy = \frac{\Gamma(\alpha+1)}{B(\alpha+2,\beta+2)\beta^{\alpha+2}} \left\|\phi'\right\|^2_{\mu_{(\alpha+1)}}.
\end{aligned}
$$

Then,

$$
\begin{aligned}
\frac{1}{\Gamma(\alpha+1)} \int_0^k |H_{\beta,k}^N(y)|^2 y^{\alpha} e^{-y} dy &\le \frac{\beta^{\alpha+2} B(\alpha+2,\beta+2)}{\Gamma(\alpha+1)N} \frac{\Gamma(\alpha+1)}{B(\alpha+2,\beta+2)\beta^{\alpha+2}} \left\|\phi'\right\|^2_{\mu_{(\alpha+1)}} \\
&= \frac{1}{N} \left\|\phi'\right\|^2_{\mu_{(\alpha+1)}},
\end{aligned}
$$

hence,

$$
\frac{1}{\Gamma(\alpha+1)} \int_0^{\infty} |H_{\beta,k}^N(y)|^2 y^{\alpha} e^{-y} dy \le \frac{C}{N} \left\|\phi'\right\|^2_{\mu_{(\alpha+1)}}.
$$

Therefore, again by Bourbaki-Alaoglu's theorem, there exists a sequence $\{\beta_j\}_{j\in\mathbb{N}}$ with $\lim_{j\to\infty} \beta_j = \infty$ such that, for all $N \in \mathbb{N}$, $\{H^N_{\beta_j,k}\}_{j\in\mathbb{N}}$ converges weakly in $L^2\left((0,\infty),\mu_\alpha\right)$ to a function $H_k^N \in L^2\left((0,\infty),\mu_\alpha\right)$. Moreover,

$$
\int_{-\infty}^{\infty} \left|H_k^N(y)\right|^2 y^{\alpha} \frac{e^{-y}}{\Gamma(\alpha+1)} dy \le \frac{C}{N}. \tag{2.33}
$$

Then, there exists a non decreasing sequence $\{N_j\}_{j\in\mathbb{N}}$ such that

$$
H_k^{N_j}(y) \longrightarrow 0, \quad a.e.\ \ y \in \mathbb{R}. \tag{2.34}
$$

Now, defining, for each $m \in \mathbb{N}$, $F_k^{N_m} = F - H_k^{N_m}$, then since $F_{\beta,k} \to F_k$, as $j \to \infty$, in the weak topology of $L^2\left((0,\infty),\mu_\alpha\right)$ and that$F^N_{\beta,k} = F_{\beta,k} - H^N_{\beta,k}$, we have that for all $m \in \mathbb{N}$,

$$
F^{N_m}_{\beta_j,k} \longrightarrow F_k^{N_m}, \quad \text{weakly in } \ L^2\left((0,\infty),\mu_\alpha\right) \quad \text{as} \quad j \to \infty. \tag{2.35}
$$

Additionally, from (2.34) we have

$$F_k^{N_m}(y) \longrightarrow F(y), \ \text{ as } m \to \infty, \ \ a.e. \ \ y \in \mathbb{R}. \tag{2.36}$$

To finish the proof of ii) we need to prove that F is the Laguerre-Riesz transform of ϕ, i.e.

$$F(y) = -\sum_{n=1}^{\infty} \frac{n!\Gamma(\alpha+1)}{(n+\alpha+1)} \widehat{\phi}^{\alpha}(n) \frac{y^{1/2}}{n^{1/2}} L_{n-1}^{\alpha+1}(y). \tag{2.37}$$

From the asymptotic relation between Jacobi and Laguerre polynomials (1.34), it is easy to see that

$$\lim_{\beta\to\infty} P_{n-1}^{(\alpha+1,\beta+1)}\left(1-\frac{2y}{\beta}\right) \ = \ \lim_{\beta\to\infty} P_{n-1}^{(\alpha+1,\beta+1)}\left(1-\frac{2y}{\beta+1}\frac{(\beta+1)}{\beta}\right) = L_{n-1}^{\alpha+1}(y).$$

On the other hand,

$$\lim_{\beta\to\infty} \widehat{\phi_\beta}(n) \ = \ \frac{1}{\Gamma(\alpha+1)} \int_0^\infty \phi(y) L_n^{\alpha}(y) y^{\alpha} e^{-y} dy = \widehat{\phi}^{\alpha}(n),$$

and

$$\begin{aligned} \lim_{\beta\to\infty} \frac{1}{\hat{h}_n^{(\alpha,\beta)}} \ &= \ \lim_{\beta\to\infty} (\frac{2n}{\beta} + \frac{\alpha}{\beta} + 1 + \frac{1}{\beta}) n! \frac{\Gamma(\alpha+1)}{(n+\alpha+1)} \frac{\beta^{\alpha+1}\Gamma(\beta+1)}{\Gamma(\alpha+\beta+2)} \frac{\Gamma\left(\beta+n+\alpha+1\right)}{\beta^{\alpha}\Gamma\left(n+\beta+1\right)} \\ &= \ \frac{n!\Gamma(\alpha+1)}{(n+\alpha+1)}. \end{aligned}$$

Then, for $y \in (0,k)$

$$\begin{aligned} \lim_{\beta\to\infty} F_{\beta,k}^{N}(y) \ &= \ -\lim_{\beta\to\infty} \sum_{n=1}^{N} \frac{\widehat{\phi_\beta}(n)}{\hat{h}_n^{(\alpha,\beta)}} \frac{(n+\alpha+\beta+1)^{1/2}}{n^{1/2}} P_{n-1}^{(\alpha+1,\beta+1)}\left(1-\frac{2}{\beta}y\right) \\ &\qquad\qquad \times \left(\frac{y}{\beta}\right)^{1/2} \left(1-\frac{y}{\beta}\right)^{1/2+\beta/2} e^{\frac{y}{2}} \\ &= \ -\lim_{\beta\to\infty} \sum_{n=1}^{N} \frac{\widehat{\phi_\beta}(n)}{\hat{h}_n^{(\alpha,\beta)}} \frac{(\frac{n}{\beta}+\frac{\alpha}{\beta}+1+\frac{1}{\beta})^{1/2}}{n^{1/2}} P_{n-1}^{(\alpha+1,\beta+1)}\left(1-\frac{2}{\beta}y\right) \\ &\qquad\qquad \times y^{1/2} \left(1-\frac{y}{\beta}\right)^{1/2+\beta/2} e^{\frac{y}{2}} \\ &= \ -\sum_{n=1}^{N} \frac{n!\Gamma(\alpha+1)}{(n+\alpha+1)} \widehat{\phi}^{\alpha}(n) \frac{y^{1/2}}{n^{1/2}} L_{n-1}^{\alpha+1}(y). \end{aligned}$$

Thus,

$$F_k^{N_m} = -\sum_{n=1}^{N_m} \frac{n!\Gamma(\alpha+1)}{(n+\alpha+1)} \widehat{\phi}^{\alpha}(n) \frac{y^{1/2}}{n^{1/2}} L_{n-1}^{\alpha+1}(y), \tag{2.38}$$

and therefore we get (2.37),

$$F(y) = -\sum_{n=1}^{\infty} \frac{n!\Gamma(\alpha+1)}{(n+\alpha+1)} \widehat{\phi}^{\alpha}(n) \frac{y^{1/2}}{n^{1/2}} L_{n-1}^{\alpha+1}(y). \ \blacksquare$$

REFERENCES

[1] Andrews, G. E., Askey, R. & Roy, R. *Special Functions* Enciclopedia of Mathematics and its Applications. Vol. 7. Cambridge University Press (2000).

[2] Bayley, W. N. *Generalized hypergeometric series.* Cambridge Tracs in Mathematics and Mathematical Physics # 32. New York. (1964).

[3] Betancor, J. Fariña, J. Rodriguez,L. Sanabria, A. *Transferring boundedness from conjugate operators associated whit Jacobi, Laguerre and Fourier-Bessel expansions to conjugate operators in the Hankel setting.* J. Fourier Anal. Appl. 14 (2008), no. 4, 493–513.

[4] Cafarelli, L. *Sobre la conjugación y sumabilidad de series de Jacobi.* Ph.D. Thesis Universidad de Buenos Aires, Argentina (1971).

[5] Gasper, G. Unpublished manuscrip.

[6] Gundy, R. *Sur les Transformations de Riesz pour le semigroup d'Ornstein-Uhlenbeck.* C.R. Acad. Sci. 303 (série I) (1986) 967-970.

[7] Gutiérrez, C. *On the Riesz transforms for the Gaussian measure.* J. Func. Anal. 120 (1) (1994) 107-134.

[8] Li, Zh.-K. *Conjugated Jacobi Series and Conjugated Functions*, J. Approx Theory, 86,(1996) 179-196.

[9] Meyer, P. A. *Quelques résultats analytiques sur le semígruope d'Ornstein-Uhlenbeck en dimension infinie.* Lectures Notes in Contr. and Inform. Sci. 49Springer-Verlag. (1983). 201-214.

[10] Meyer, P. A. *Transformations de Riesz pour les lois Gaussiennes.* Lectures Notes in Math 1059 Springer-Verlag. (1984). 179-193.

[11] Muckenhoupt, B. *Hermite conjugate expansions.* Trans. Amer. Math. Soc. 139 (1969) 243-260.

[12] Muckenhoupt, B. *Conjugate functions for Laguerre expansions.* Trans. Amer. Math. Soc. 147 (1970) 403-418.

[13] Muckenhoupt, B. & Stein, E.M. *Classical Expansions.* Trans. Amer. Math. Soc. 147 (1965) 17-92.

[14] Nowak, A. *On Riesz transforms for Laguerre expansions.* J. Funct. Anal. 215 (2004), no. 1, 217–240.

[15] Nowak, A. ; Sjögren, P. *Riesz transforms for Jacobi expansions.* J. Anal. Math. 104 (2008), 341–369.

[16] Nowak, A. ; Stempak, K. *L^2-theory of Riesz transforms for orthogonal expansions.* J. Fourier Anal. Appl. 12 (2006), no. 6, 675–711.

[17] Pérez, S.; Soria, Fernando *Operators associated with the Ornstein-Uhlenbeck semigroup.* J. London Math. Soc. (2) 61 (2000), no. 3, 857–871.

[18] Pisier, G. *Riesz Transform: a simpler analytic proof of P. A. Meyer inequality.* Lectures Notes in Math 1321. Springer-Verlag (1988) 485-501.

[19] Sjögren, P. *Operators associated with the Hermite semigroup-a survey.* J. Fourier Anal. Appl. 3 (1997), Special Issue, 813–823.

[20] Stein, E.M. *Singular Integrals and differenciability properties of functions.* Princeton Univ. Press. Princeton (1970) .

[21] Stein, E.M. *Topics in Harmonic Analysis related to the Littlewood-Paley Theory.* Princeton Univ. Press. Princeton (1970).

[22] Szegö, G. *Orthogonal polynomials.* Colloq. Publ. 23. Amer. Math. Soc. Providence (1959).

[23] Urbina, W. *Singular Integrals with respect to the Gaussian measure.* Scuola Normale Superiore di Pisa. Classe di Science. Serie IV Vol XVIII, 4 (1990) 531-567.

[24] Urbina, W. *Operators Semigroups associated to Classical Orthogonal Polynomials and Functional Inequalities.* Seminaires et Congres. French Mathematical Society (SMF). 200812.

[25] Zygmund, A. *Trigonometric Series.* 2nd. ed. Cambridge Univ. Press. Cambridge (1959).

Departmento de Matemáticas, Universidad Experimental Francisco de Miranda, Punto Fijo, Venezuela.
E-mail address: `[Eduard Navas]eduard.navas@gmail.com`

Department of Mathematical and Actuarial Sciences, Roosevelt University Chicago, Il, 60605, USA.
E-mail address: `[Wilfredo Urbina]wurbinaromero@roosevelt.edu`